

Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =
Bulletin suisse de minéralogie et pétrographie

Band: 66 (1986)

Heft: 3

Artikel: Trabzonite $\text{Ca}_4\text{Si}_3\text{O}_{10} \cdot 2\text{H}_2\text{O}$ a new hydrated calcium silicate

Autor: Sarp, Halil / Burri, Georges

DOI: <https://doi.org/10.5169/seals-50903>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 13.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Trabzonite $\text{Ca}_4\text{Si}_3\text{O}_{10} \cdot 2 \text{H}_2\text{O}$ a new hydrated calcium silicate

by *Halil Sarp*¹ and *Georges Burri*²

Abstract

Trabzonite, ideally $\text{Ca}_4\text{Si}_3\text{O}_{10} \cdot 2 \text{H}_2\text{O}$, has been found near Ikizdere at NW of Varda Yaylasi (Rize) (near Trabzon county) - Turkey and occurs in skarn produced by granitic intrusion in volcano-sedimentary formations. Associated minerals are: spurrite, rustumite, perovskite, calcite, vesuvianite, tobermorite, garnet, killalaite, hillebrandite and molybdenite. The crystals, colourless, are small irregular or hypidiomorphs from 0.1 to 0.2 mm with a white streak and vitreous lustre; transparent. The hardness could not be measured because of small size. Chemical composition determined by electron microprobe is: CaO 50.2, MgO 0.13, MnO 0.04, Na₂O 0.3; Al₂O₃ 0.05, SiO₂ 40.38 and H₂O 7 (weight loss), total 98.1%. The crystal system is monoclinic, space group P_{2_1} or P_{2_1}/m with $a = 6.895$ (2), $b = 20.640$ (3), $c = 6.920$ (2) Å, $\beta = 98^\circ$ and $Z = 4$. The $a:b:c$ ratio calculated from unit-cell parameters is 0.3341:1:0.3353. The strongest lines in the X-ray powder diffraction pattern (dÅ for CuK α , Ivis., hkl) are: 5.71 (35) (021, 120), 3.442 (60) (060, 002), 3.062 (100) (032, 230), 2.912 (30) (102, 201), 2.851 (50) (042, 240), 2.635 (50) (052, 250) and 2.585 (90) (212, 080). The density is 2.9 (meas.) and 3.08 g/cm³ (calc.). Optically, Trabzonite is biaxial (+) with $2V$ (meas.) = 55(5)°, $2V$ (calc.) = 60°; $\alpha = 1.632$ (2), $\beta = 1.634$ (2), $\gamma = 1.640$ (2) (589 nm). Dispersion $r > v$ weak to moderate. Optical orientation: $X \wedge c = 8^\circ$, $Y = b$, $Z = a$. Trabzonite is named for to honour the vilâyet of Trabzon. This new mineral and mineral name have been approved by the Commission on New Minerals and Mineral Names of the International Mineralogical Association. The complete description of the mineral has been presented at the congress of T.J.K. at Ankara.

Keywords: New mineral, trabzonite, Turkey.

¹ Halil Sarp, Département de Minéralogie, Muséum d'Histoire naturelle de Genève, case postale 434, CH-1211 Genève 6, Suisse.

² Georges Burri, Institut de Physique Expérimentale, Université de Lausanne, Laboratoire de Microsonde, CH-1015 Lausanne-Dorigny, Suisse.