

Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =
Bulletin suisse de minéralogie et pétrographie

Band: 71 (1991)

Heft: 3

Artikel: Geochemistry and tectonic significance of Late Hercynian potassic and
ultrapotassic magmatism in the Aar Massif (Central Alps)

Autor: Schaltegger, Urs / Gnos, Edwin / Küpfer, Theo

DOI: <https://doi.org/10.5169/seals-54372>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 13.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Geochemistry and tectonic significance of Late Hercynian potassic and ultrapotassic magmatism in the Aar Massif (Central Alps)

by Urs Schaltegger¹, Edwin Gnos², Theo Küpfer³ and Toni P. Labhart²

Abstract

In the eastern Aar Massif (Central Alps, Switzerland) two different Upper Carboniferous intrusive series can be recognized: (A) A shoshonitic-ultrapotassic series, consisting of dioritic rocks of Val Punteglias, of the Giuv syenite and the Punteglias granite, and (B) a high-K calc-alkaline series consisting of the Düssi-Fruttstock diorite and the Brunni granite, both situated in the Brunnital. Both series originate from an enriched mantle source high in Rb, K and Th, but experienced different processes during their evolution. The shoshonitic-ultrapotassic series was possibly contaminated by partial melts of subducted slab sediments, but was hardly affected by fractionation processes; it therefore represents rather original material from the subcontinental mantle. The calc-alkaline series experienced extensive fractional crystallization and contamination by crustal melts.

The shoshonitic-ultrapotassic rocks were emplaced at a late stage of Hercynian orogeny. They were intruded after the cessation of compression and mark the beginning of late Hercynian strike-slip tectonics that enabled mantle melts to rise into the upper crust.

Keywords: ultrapotassic rocks, geochemistry, mantle-crust interaction, Hercynian magmatism, Aar Massif, Central Alps, Switzerland.

Introduction

Potassic and ultrapotassic rocks are rare but occur in various plate tectonic settings; they may be part of continental rift systems or intrude into regions of post-orogenic crustal extension and heating (e.g. THOMPSON et al., 1989). Shoshonitic to ultrapotassic rocks are widespread among ancient and modern volcanic and magmatic arcs, e.g. in the Eolian arc (ELLAM et al., 1989), Sunda-Banda arc (WHELLER et al., 1987), late tectonic plutons in the Grenville Province (CORRIVEAU, 1990) and also in Archean granite-greenstone terrains (SHIREY and HANSON, 1984). Shoshonitic to ultrapotassic rocks may also occur in orogenic settings, along active continental margins (THOMPSON and FOWLER, 1986).

Genetic models for shoshonitic-ultrapotassic rock associations often imply the existence of a LILE (large ion lithophile element)-enriched subcontinental mantle source; variable crustal contamination and different extents of fractional crystallization in intermediate and final magma chambers lead to a large geochemical diversity of different shoshonitic rock suites. The geotectonic setting is generally thought to be distant from the active subduction, thus the melts are of deep-seated mantle origin and were generated and fractionated in either hydrous or anhydrous environments.

We report here the occurrence of a potassic to ultrapotassic magmatic association of late Hercynian age, situated in the pre-Mesozoic basement of

¹ Centre de Géochimie de la Surface, CNRS, 1, rue Blessig, F-67084 Strasbourg, France.

² Mineralogisch-petrographisches Institut, Universität Bern, Baltzerstrasse 1, 3012 Bern, Switzerland.

³ Nationale Gesellschaft für die Lagerung radioaktiver Abfälle (NAGRA), Parkstrasse 23, 5401 Baden, Switzerland.

the Central Alps. The aim of this paper is to describe the petrography, the geological setting and the geochemistry of these rocks and to summarize the current ideas about the genesis of these rocks as a basis for future work.

Geology

The Aar Massif is part of the pre-Mesozoic basement in the external zone of the Hercynian fold belt (Helvetic zone, FRISCH *et al.*, 1990). The Helvetic nappes above the massif were removed during Alpine and post-Alpine uplift and erosion. The massif consists of Hercynian magmatic and volcanic rocks that were intruded into a polymetamorphic basement of pre-Hercynian age (Fig. 1). The late Hercynian intrusives contain granites, diorites, monzonites and syenites and were emplaced 334 to 298 Ma ago (SCHALTEGGER *et al.*, 1991). Volcanic and volcanoclastic sequences occur throughout the whole massif (BÖHM, 1988; FRANKS, 1968 a, b; SCHENKER, 1986; MERCOLLI *et al.*, 1988); field evidence and geochemical criteria are consistent with a similar Upper Carboniferous extrusion/deposition age for the volcanic rocks. The late Hercy-

nian intrusive rocks can be subdivided into three different groups, according to geochemical signature and age:

(A) A shoshonitic to ultrapotassic suite (SUPS) in the eastern part of the massif, comprising the Giv syenite (HUBER, 1948; LABHART and RYBACH, 1971; LABHART, 1977) and the diorites and granites of Val Punteglias (KÜPFER, 1977; SEEMANN, 1975), emplaced 334 Ma ago (SCHALTEGGER *et al.*, 1991).

(B) A high-K calc-alkaline series (KCAS), comprising the diorite-granite complex of Brunnital (Düssi-Fruttstock diorite and Brunni granite; GNOS, 1988, SIGRIST, 1947) and various other dioritic to granodioritic rocks in the central part of the massif, altogether emplaced around 308–310 Ma (SCHALTEGGER *et al.*, 1991).

(C) The highly differentiated Central Aar granite, the largest late Hercynian intrusion, was emplaced 298 Ma ago (SCHALTEGGER and VON QUADT, 1990; SCHALTEGGER *et al.*, 1991). Its petrography and geochemistry is extensively described in SCHALTEGGER (1989, 1990) and thus will not be discussed in detail in this paper. Data of SCHALTEGGER (1990) are however included in some diagrams to allow direct comparison of all three magmatic stages.




Fig. 1 Geologic sketch map of the central and eastern parts of the Aar Massif. Symbols: A) Sedimentary cover of the Aar Massif; B) Tscharren-Formation, volcanoclastic rocks; C) Val Gliems-Formation, volcanoclastic rocks; D) Central Aar granite; E) Southern Aar granite; F) Brunni granite; G) Düssi-Fruttstock diorite; H) Punteglias diorite; I) Punteglias granite; K) Giv syenite; L) Polymetamorphic basement; M) Southern limit of the Aar Massif.

All rocks suffered a greenschist facies metamorphic overprint of Alpine age (FREY et al., 1980) that caused various mineral reactions and brittle to plastic deformation. This overprint modified geochemical compositions along shear zones (MARQUER, 1987) and disturbed or reset isotopic mineral systems (DEMPSTER, 1986). Late-Hercynian deformation and formation of mylonites must be envisaged, too.

Petrography of the investigated rocks

SHOSHONITIC-ULTRAPOTASSIC SERIES

This group includes three different rock units: The *diorites to monzodiorites of Val Punteglia*s range from meladiorites and biotite-hornblendites to quartz-monzonites and quartz-monzodiorites in composition (Fig. 2). They consist of 8 to 10 small intrusive stocks, covering a total surface area of less than 1 km². The more mafic rock types are coarse-grained and contain up to 90% (by volume) of mafic minerals, the abundance of hornblende always exceeding biotite. The biotite-hornblende textures are considered to be magmatic. The mineral assemblage always contains normative ol. The more leucocratic quartz-monzonites and monzodiorites are fine-grained and inhomogeneous. On a mesoscopic scale, igneous layering is widespread in the monzosyenitic to dioritic rocks; the distribution of hornblende and biotite cumulates and plagioclase phenocrysts is patchy. The members of this group contain anhedral K-feldspar and euhedral acicular apatite, allanite, titanite, epidote, ilmenite, pyrite and zircon as accessory phases. The field relationships are typical for in-situ magma mixing: transitional contacts, layering with alternating crystal-rich horizons, magmatic flow textures and dioritic globules in a monzosyenitic matrix. Dykes crosscutting these rocks are generally more differentiated.

The *Punteglia granite* occurs in the same region, intruding into the diorites and suggesting a slightly younger age. It forms a lense-shaped body covering an area of 10 km² (Fig. 1). The granite ranges from a monzogranitic to granodioritic composition and can be subdivided in melanocratic, normal and leucocratic facies, respectively, and is rimmed and intruded by small stocks and dykes of a genetically related leucogranite, the Postabiala granite (SEEMANN, 1975). In the QAP diagram (Fig. 2) the granitic association reveals an increasing plagioclase/K-feldspar ratio in the very latest steps of differentiation. The granite contains large amounts of K-feldspar phenocrysts forming magmatic flow




Fig. 2 QAP diagram showing the variation of modal compositions of the investigated rocks. Data from GNOS (1988), LABHART (1977), LABHART and RYBACH (1971), SEEMANN (1975); the two values for the Punteglia diorites are mean values of KÜPFER (1977) for a more mafic (meladiorite to biotite-hornblendite) and an intermediate group (quartz-monzonite to quartz-monzodiorite).

textures in a matrix of quartz, partly euhedral plagioclase (An 5–20), biotite and euhedral hornblende. Accessory minerals include apatite, titanite, epidote, allanite, zircon and opaques.

The *Gjuv syenite* is the ultrapotassic member of this group; it occurs as a small lens (6 km²) some 20 km west of Val Punteglia and has no visible connection with the above discussed rocks. It consists of a porphyritic coarse-grained central and a finer-grained marginal facies, both containing large euhedral K-feldspar phenocrysts showing magmatic flow textures. A second K-feldspar generation rims the phenocrysts and forms anhedral crystals in a matrix consisting of quartz, plagioclase and patches of biotite and hornblende. Accessories are titanite, apatite, zircon, allanite, epidote and opaques. Modal analyses plot in the fields of quartz-monzonite and quartz-syenite of a QAP diagram (Fig. 2). The normative mineralogy is partly SiO₂-undersaturated, few have normative ol.

HIGH-K CALC-ALKALINE SERIES

This group includes the leucocratic Brunni granite and an association of diorites in the region of Brunntal (Düssi-Fruttstock; Fig. 1). The *diorites* range from cumulate-like hornblende-gabbros to hornblende-diorites and hornblende- or biotite-quartz-monzonites. This wide range of intermediate rock composition can coexist at the scale of meters to decameters. The more mafic rock types typically occur as inclusions in more acid parts, sometimes forming magmatic breccia but more often having transitional contacts. These field characteristics suggest the contemporaneous intrusion of all rock types. Sometimes, they also contain angular xenoliths of biotite-plagioclase-gneiss or amphibolite from the adjacent basement. In more mafic rocks euhedral brown titaniferous hornblende cores are overgrown by green hornblende and titanite. Euhedral apatite and allanite/epidote occur as early liquidus phases, anhedral quartz, K-feldspar and plagioclase are late crystallizing phases. In some quartz-bearing rocks skeletal zircon was observed. Along the border of the intrusion, quartz-bearing diorites containing skeletal hornblende are common. Locally they are associated with pegmatoid pockets or dioritic globules in a syenitic matrix; towards the Brunni granite, K-feldspar phenocrysts are abundant.

The *Brunni granite* is a leucogranite containing 2–3 vol.% of mafic minerals. Large crystals of allanite/epidote are very characteristic. In a few places a border facies with K-feldspar megacrysts was observed. The granite is very homogeneous and shows a schistosity, presumably of Alpine age, which is more intense towards the south. An age of 308 ± 2 Ma was determined by SCHALTEGGER et al. (1991) from a diorite of this intrusive complex.

One sample of the *Schöllenen diorite* (Reuss valley, KAW 2520) has also been included in this study. This rock is coeval with the Düssi-Fruttstock diorite (SCHALTEGGER et al., 1991) and occurs as an enclave within the Central Aar granite.

Geochemistry

ANALYTICAL PROCEDURES

Geochemical analyses of the rocks from Val Puntiglias are taken from KÜPFER (1977). Major and trace elements were analyzed by X-ray fluorescence (model Philips PW 1410) without specific matrix correction. Analytical precision is 50% concentration per weight $\pm 2\%$ (relative), $1\% \pm 4\%$, 1000 ppm $\pm 6\%$, 50 ppm $\pm 15\%$ (1 sigma standard deviations). K, Na, FeO, H₂O and CO₂ were determined using wet chemical methods.




Fig. 3 K₂O vs SiO₂ (% per weight) diagram showing the classification of PECERILLO and TAYLOR (1976). Same symbols as in figure 2. The rocks of the shoshonitic-ultrapotassic suite (SUPS, filled symbols) lie in the fields of absarokite (ab), shoshonite (sh) and banakite (ba), the high-K, calc-alkaline rocks (KCAS) in the fields of absarokite, shoshonite, high-K andesite (ka) and dacite (da). Granites plot into the fields of high-K dacite (kd) and rhyolite (rh). The data for the Puntiglias granite are mean values for different granite facies from SEEMANN (1975); Central Aar granite data from SCHALTEGGER (1989).

The geochemical data from the diorite-granite complex of the Brunntal are taken from GNOS (1988). Major elements were measured on fused rock tablets, trace elements on powder pills by XRF on a PW 1450 Philips spectrometer. Analytical errors are 50% $\pm 1\%$, 1% $\pm 5\%$, 1000 ppm $\pm 5\%$, 100 ppm $\pm 10\%$, 10 ppm $\pm 20\%$ (1 sigma). FeO, H₂O and CO₂ were determined using wet chemical methods.

The Giuv syenite data were determined in 1989 at the Centre d'Analyse Minérale de l'Université de Lausanne on a Philips PW 1400 X-ray fluorescence spectrometer. Fused tablets were used for major elements, pressed powder pills for trace elements. Analytical errors are 50% $\pm 1\%$, 1000 ppm $\pm 5\%$, 100 ppm $\pm 10\%$ (1 sigma). FeO, H₂O and CO₂ were determined using wet chemical methods.

The rare earth element analyses were carried out by Bondar-Clegg & Company Ltd. (Ottawa, Canada). Informations on analytical precision are not available. Detection limits are: La 0.5 ppm, Ce 5 ppm, Sm 0.1 ppm, Nd 10 ppm, Eu 1 ppm, Dy 1 ppm, Yb 0.5 ppm, Lu 0.1 ppm.

MAJOR AND TRACE ELEMENTS

The geochemical compositions for some representative samples are listed in table 1. The investigated rocks can be divided into two groups according to their geochemical characteristics:

Tab. 1 Major and trace element concentration of selected samples: Numbers 1 to 11 shoshonitic-ultrapotassic series (SUPS), 12 to 17 high-K, calc-alkaline series (KCAS). 1–4 Punteglias granite association (average values from SEEMANN, 1975): 1 melanocratic, 2 normal, 3 leucocratic facies, 4 Postabiala granite; 5–9 Giuv syenite: 5 coarse-grained (KAW 252), 6 enclave in the Central Aar granite (KAW 253), 7 and 8 both fine-grained (KAW 254 and 255), 9 coarse-grained (GS 23); 10–11 diorites Val Punteglias: 10 biotite-hornblende (TK 196), 11 quartz-monzonite (TK 284), from KÜPPER (1977); 12–14 Düssi diorite: 12 hornblende diorite (EG 60), 13 biotite-hornblende diorite (EG 124), 14 biotite-hornblende tonalite (EG 126); 15–16 Brunni granite: leucogranites (EG 139 and 147), from GNOS (1988); 17 Schöllenen diorite: biotite-hornblende diorite (from SCHALTEGGER, 1989). Complete data set of the Giuv syenite analyses may be obtained on request from the first author.

element	SUPS																	KCAS																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
	Punteglias granite			Giuv syenite														Punteglias diorite			Düssi diorite				Brunni granite			Schöllenen diorite																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000	1001	1002	1003	1004	1005	1006	1007	1008	1009	1010	1011	1012	1013	1014	1015	1016	1017	1018	1019	1020	1021	1022	1023	1024	1025	1026	1027	1028	1029	1030	1031	1032	1033	1034	1035	1036	1037	1038	1039	1040	1041	1042	1043	1044	1045	1046	1047	1048	1049	1050	1051	1052	1053	1054	1055	1056	1057	1058	1059	1060	1061	1062	1063	1064	1065	1066	1067	1068	1069	1070	1071	1072	1073	1074	1075	1076	1077	1078	1079	1080	1081	1082	1083	1084	1085	1086	1087	1088	1089	1090	1091	1092	1093	1094	1095	1096	1097	1098	1099	1100	1101	1102	1103	1104	1105	1106	1107	1108	1109	1110	1111	1112	1113	1114	1115	1116	1117	1118	1119	1120	1121	1122	1123	1124	1125	1126	1127	1128	1129	1130	1131	1132	1133	1134	1135	1136	1137	1138	1139	1140	1141	1142	1143	1144	1145	1146	1147	1148	1149	1150	1151	1152	1153	1154	1155	1156	1157	1158	1159	1160	1161	1162	1163	1164	1165	1166	1167	1168	1169	1170	1171	1172	1173	1174	1175	1176	1177	1178	1179	1180	1181	1182	1183	1184	1185	1186	1187	1188	1189	1190	1191	1192	1193	1194	1195	1196	1197	1198	1199	1200	1201	1202	1203	1204	1205	1206	1207	1208	1209	1210	1211	1212	1213	1214	1215	1216	1217	1218	1219	1220	1221	1222	1223	1224	1225	1226	1227	1228	1229	1230	1231	1232	1233	1234	1235	1236	1237	1238	1239	1240	1241	1242	1243	1244	1245	1246	1247	1248	1249	1250	1251	1252	1253	1254	1255	1256	1257	1258	1259	1260	1261	1262	1263	1264	1265	1266	1267	1268	1269	1270	1271	1272	1273	1274	1275	1276	1277	1278	1279	1280	1281	1282	1283	1284	1285	1286	1287	1288	1289	1290	1291	1292	1293	1294	1295	1296	1297	1298	1299	1300	1301	1302	1303	1304	1305	1306	1307	1308	1309	1310	1311	1312	1313	1314	1315	1316	1317	1318	1319	1320	1321	1322	1323	1324	1325	1326	1327	1328	1329	1330	1331	1332	1333	1334	1335	1336	1337	1338	1339	1340	1341	1342	1343	1344	1345	1346	1347	1348	1349	1350	1351	1352	1353	1354	1355	1356	1357	1358	1359	1360	1361	1362	1363	1364	1365	1366	1367	1368	1369	1370	1371	1372	1373	1374	1375	1376	1377	1378	1379	1380	1381	1382	1383	1384	1385	1386	1387	1388	1389	1390	1391	1392	1393	1394	1395	1396	1397	1398	1399	1400	1401	1402	1403	1404	1405	1406	1407	1408	1409	1410	1411	1412	1413	1414	1415	1416	1417	1418	1419	1420	1421	1422	1423

(A) The shoshonitic-ultrapotassic series (SUPS) shows an increase in SiO_2 by 17%, in K_2O by 41% and a decrease in MgO by 71% from the most mafic to the most felsic samples. In the classification scheme of PECERILLO and TAYLOR (1976), which was established for volcanic rocks, most data points are situated in the absarokite, shoshonite and banakite fields, the granitic rocks plot into the high-K dacite and rhyolite fields (Fig. 3). The Givv syenite is the only member that meets the requirements of the classification of FOLEY et al. (1987) for ultrapotassic rocks ($\text{K}_2\text{O}/\text{Na}_2\text{O} > 2\%$, $\text{K}_2\text{O} > 3\%$, $\text{MgO} > 3\%$).

The spidergram (chondrite-normalized variation diagram in order of increasing compatibility, THOMPSON, 1982; Fig. 4) of the SUPS reveals a LILE-enrichment in the Givv syenite (samples KAW 252–255) and Punteglias granite (KAW 676), up to 3000 times chondritic values. An average diorite (major and trace elements: mean of 3 dioritic to monzodioritic samples from KÜPFER, 1977, REE of SU-89-10) and a biotite-hornblendite (TK 196) are considerably less enriched; the latter differs from the other rocks of the SUPS mainly by the smaller presence of Ba, Rb, Sr and LREE. Troughs at Nb (Ta), Sr and Ti are generally taken as indicators of a genesis by subduction-related processes (THOMPSON et al., 1984). For highly evolved rocks like the discussed syenites and granites, these troughs are astonishingly weak (see syenites of THOMPSON and FOWLER, 1986, for comparison). Compared to lamprophyres that are coeval with group C granites (OBERHÄNSLI, 1986) the SUPS rocks show stronger LILE-enrichment and less depleted Nb, P, Sr and Ti. According to their high Al_2O_3 , CaO and relatively low MgO and TiO_2 contents, the ultrapotassic rocks are classified into group III of FOLEY et al. (1987), typical for orogenic ultrapotassic rocks. They, however, have higher LILE and lower contents on the more compatible end of the element selection (La to Yb) as well as less marked troughs for Nb, Sr and Ti, compared to group III standard rocks of FOLEY et al. (1987).

(B) The high-K calc-alkaline series (KCAS) starts with rocks of shoshonitic composition, too, but evolves towards lower potassium contents with increasing acidity (Fig. 3). It contains no ultrapotassic members. The leucogranitic counterpart, the Brunni granite, has K_2O contents around 5%, which are normal for high-Si melts. The spidergrams of the KCAS (Fig. 4) show a much smaller LILE-enrichment and are more irregular. The diorites exhibit large variations in La, Ce, Sr, P and Zr, whereas Nb and Ti are relatively consistent (Fig. 4), reflecting either variable allanite, plagioclase, apatite and zircon fractionation or mixing of melts with different compositions. Allanite certainly was an




Fig. 4 Spidergrams (chondrite-normalized variation diagrams in increasing compatibility order, after THOMPSON, 1982; normalizing factors for Rb, K and P after SUN, 1980); shoshonitic-ultrapotassic series (SUPS): Givv syenite (KAW 252–255), Punteglias granite (KAW 676) and diorites of Val Punteglias (SU-89-10, TK 196); high-K calc-alkaline series (KCAS): diorites of Düssi-Frutstock complex (EG 61–183) and Schöllenen diorite (KAW 2520); and Brunni granite: leucogranites (EG 139, 141, 147, 162).

important fractionating mineral, judging from the highly variable Th, La and Ce concentrations. The Brunni leucogranite spidergram (Fig. 4) has more marked negative peaks at Ba, Th, Nb, P and Ti. The Ba depletion reflects K-feldspar crystallization in the latest stage of differentiation (MITTFELDT and MILLER, 1983; SCHALTEGGER and KRÄHENBÜHL, 1990).

The Central Aar granite shows an evolution from the high-K dacite to rhyolite fields in the K_2O vs SiO_2 classification screen and thus follows a trend which is again different from the ones of the SUPS and KCAS.




Fig. 5 AFM diagram. Same symbols as in figure 2.

The rocks of both SUPS and KCAS plot on a low-Fe calc-alkaline trend in a AFM diagram (Fig. 5) which is typical for shoshonitic series (MORRISON, 1980). The selected Harker diagrams (Fig. 6 a, b) show the compatible behaviour of Mg and Ti (similar relationship for P, Ca and Fe) and do not discriminate between the SUPS and KCAS. Smooth fractionation trends for higher silica rocks (Giv. syenite; Brunni, Punteglias and Central Aar




Fig. 6 Harker diagrams; a: MgO, b: TiO₂. Same symbols as in figure 2.




Fig. 7 Zr vs SiO₂ diagram. Same symbols as in figure 2.




Fig. 8 Sr vs Rb diagram. Same symbols as in figure 2.

granites) contrast with the data scatter observed for the diorites, indicating the involvement of an additional process such as mixing of different melts or source components. K₂O, Na₂O, Th and U behave mostly incompatibly during the evolution of the SUPS, elements like Rb, Sr or Zr change to compatible geochemical behaviour at certain SiO₂ contents (Figs 7 and 8): Zr reaches saturation in the most silicic dioritic rocks; the syenite is saturated and shows a differentiation into two groups with higher and lower Zr contents, respectively (Fig. 7). The behaviour of Sr is similar: a low-Sr and high-Sr series can clearly be recognized (Fig. 8). They are not distinguishable on petrographical grounds and do not represent the separation into central and border facies at all. The Zr-SiO₂ systematic of the KCAS diorites is difficult to understand, but probably follows a similar evolutionary path as the SUPS diorites (Fig. 7); Rb and Sr, however, show a negative correlation (Fig. 8). Few outliers deviate from the trend lines in figures 7 and 8, which might be interpreted with mixing and/or cumulation processes.

The transition elements Mg, Cr and Ni are closely correlated for both series (Fig. 9 a, b), indi-




Fig. 9 a: Cr vs Ni; b: MgO vs Ni. Same symbols as in figure 2. Qualitative fractionation paths for olivine and clinopyroxene are indicated. Evolutionary trend shows a hiatus at 200 ppm Cr, 70 ppm Ni and 6% MgO, indicating a change from predominantly olivine to clinopyroxene-dominated fractionation at an early stage of magma evolution.

cating a two-stage evolution. It may be interpreted as a change in the ratio of olivine to pyroxene fractionation during the crystallization in a primary magma reservoir, or it may be created by another process than fractional crystallization (e.g. mixing), which would also explain the scatter of the dioritic rocks in figures 9 a and b.

RARE EARTH ELEMENTS

The rare earth element (REE) patterns of 8 samples, 6 from the SUPS and 2 from the KCAS, reveal a moderate enrichment of light rare earth elements (LREE; Fig. 10, Tab. 2). This feature is a characteristic of potassic rocks; similar or higher LREE contents can be found in modern ultrapotassic island arc volcanics (WHELLER et al., 1987). La_N/Yb_N values are highest in the Giv syenite (24–28), lower in the Punteglias and Düssi diorites (18–20) and lowest in the Schöllenen diorite (9.6). There is little variation in the LREE concentrations; the differences are mainly caused by variable levels of heavy rare earth elements (HREE). A biotite-hornblende

dite from the SUPS (TK 196) has lower LREE concentrations suggesting lower bulk distribution coefficients for La to Sm, compared to the other samples.

The analyses do not show Eu anomalies despite the Sr variations in the spidergrams and high K-feldspar contents in the Giv syenite. The lack of an Eu anomaly will be discussed in the next chapter. The patterns are similar to continental alkali basaltic trends (CHAUVEL and JAHN, 1984), but have lower LREE and higher La_N/Yb_N ; they are smoother than patterns of the Aar Massif lamprophyres (OBERHÄNSLI et al., 1991), but show similar, slightly elevated HREE-plateaus.

Discussion

Both investigated rock suites consist mainly of dioritic and granitic rocks, intermediate syenites are only present in the SUPS and are lacking or undetected in the younger KCAS. The two series can be differentiated by means of their geochemical composition (Fig. 3). The shoshonitic-ultrapotassic series (SUPS), emplaced 334 Ma ago (SCHALTEGGER et al., 1991), is characterized by high LILE concentrations, elevated element ratios such as Ba/Sr, Ba/La, La/Yb, but low Rb/Sr and relatively primitive




Fig. 10 Rare earth element variation diagrams for the shoshonitic-ultrapotassic series (SUPS) and two samples of the high-K calc-alkaline series (KCAS). For sample number identification see caption figure 4; KAW 3120 = biotite-hornblende diorite of Düssi-Fruttstock diorite complex. Normalizing factors of NAKAMURA (1974).

Tab. 2 Rare earth element concentrations. INAA determinations from Bondar-Clegg & Company Ltd., Ottawa, Canada.

sample	rock type	La [ppm]	Ce [ppm]	Nd [ppm]	Sm [ppm]	Eu [ppm]	Tb [ppm]	Dy [ppm]	Yb [ppm]	Lu [ppm]
KAW 252	Giuv syenite, c.g.	50	112	45	9	2.5	<1.0	3	1.2	0.1
KAW 253	Giuv syenite, f.g.	43	100	45	10	2.1	<1.0	2	1.0	0.1
KAW 254	Giuv syenite, f.g.	44	96	45	6	1.8	<1.0	4	1.1	0.1
KAW 255	Giuv syenite, c.g.	50	106	51	6	2.2	<1.0	3	1.2	0.1
KAW 676	Punteglias granite	48	105	47	8	2.1	<1.0	3	1.5	0.2
KAW 2520	Schöllenen diorite	50	109	42	10	2.3	1.3	9	3.9	0.5
KAW 3120	Düssi diorite	67	120	37	8	2.0	<1.0	5	2.2	0.3
SU-89-10	Punteglias monzo- diorite	55	117	49	12	2.5	<1.0	4	2.0	0.3

abbreviations: c.g. = coarse-grained, f.g. = fine-grained

REE patterns (Fig. 10). The rocks of the high-K calc-alkaline series (KCAS) are some 25 Ma younger (SCHALTEGGER et al., 1991) and show trace element patterns that are more typical of subduction-generated magmas, indicating a strong influence of fractionation processes. They also show primitive REE-patterns (Fig. 10), but no evident LILE anomaly.

The investigated rocks of both series are late orogenic intrusions and evolve to or occur together with contemporaneous or younger calc-alkaline granitoids (SCHALTEGGER and VON QUADT, 1990; SCHALTEGGER et al., 1991). Dioritic members of both series show clear field evidences for magma mixing, which is also indicated by a late K-feldspar crystallization in diorites and syenites. Abundant magmatic flow textures were generated during the emplacement of the phenocryst-containing melts in upper crustal levels, most spectacular for the K-feldspar-rich rocks (Giuv syenite and Punteglias granite). A weak post-emplacment deformation is present in all rock types and may be of late Hercynian and Alpine age.

THE ROLE OF FRACTIONAL CRYSTALLIZATION

The rocks of the SUPS show extensive liquidus crystallization (hornblende, biotite, titanite, apatite, allanite, zircon, plagioclase and K-feldspar) indicated by the presence of phenocrysts. During fractional crystallization, Rb, K₂O and especially Th must have behaved incompatibly, leading to anomalously high contents of U and Th in the Giuv syenite (mean values of 24 and 65 ppm, respectively; see also LABHART and RYBACH, 1971). This process, however, causes only weak fractionation troughs at La, Nb, Ti or Zr (Fig. 4) and no Eu-

anomaly (Fig. 10) at all. The extent of the troughs in figure 4 does not change significantly from mafic to acid members of the SUPS, thus the low-pressure in-situ crystallization was not able to change these geochemical features. It must therefore be concluded that most of the geochemical characteristic is inherited from a common source. Any differentiation process as well as mixing with crustal components would, however, enhance the contents of Rb, Th and K. This may possibly contribute to the LILE-enrichment of the syenites of the SUPS (Fig. 4). Post-emplacment differentiation is indicated by systematic geochemical differences between different facies of the same rock type (core-rim of the Giuv syenite, see LABHART and RYBACH, 1971; melanocratic, normal and leucocratic facies of the Punteglias granite, see SEEMANN, 1975). Extensive fractional crystallization of minerals with K_D-values << 1 for LILE (pyroxenes, hornblendes, olivine) at an early magmatic stage would lead to considerable depletions in MgO, TiO₂, Cr or Ni, which is not the case for the SUPS rocks.

The rocks of the KCAS show evidence for fractional crystallization of biotite, hornblende, apatite, and eventually zircon, indicated by variable chondrite-normalized Sr, P, Ti and Zr concentrations in figure 4; largely variable La/Nb and Zr/Ti ratios may be caused by the removal of allanite and zircon. P and Ti peaks for KCAS and Brunni granite (Fig. 4) indicate that apatite fractionation is restricted to the dioritic stage and a subordinate process for the Brunni granite, whereas Ti depletion happened during the granitic stage, probably by fractionation of biotite and Ti-Fe-oxides. Compared to the SUPS fractional crystallization thus played a more important role in the genesis of the KCAS.

The importance of magma mixing processes is difficult to assess. In addition to field observations (GNOS, 1988; KÜPFER, 1977), mixing is indicated by

a large scatter in the geochemical data for the diorite samples of both series in some of the diagrams (Figs. 6 and 9). GNOS (1988) tentatively postulated a magma mixing process based on field evidence, but did not find a well fitting model.

SOURCE COMPOSITION

An assessment on compositions of parent melts remains very hypothetical without isotopic data. Compositions may be estimated from the most mafic and Mg-rich rocks of both suites. None of the investigated lithologies, however, strictly represents a pure melt composition; all of them contain phenocrysts of liquidus phases (allanite, apatite, hornblende/biotite, titanite, K-feldspar and plagioclase). The most mafic rocks of the SUPS have 14–17% MgO, 800–1200 ppm Cr and 300–600 ppm Ni; mafic analogues of the KCAS show contents of 10–15% MgO, 450–500 ppm Cr and 100–550 ppm Ni. Considering these concentrations, a genesis of both SUPS and KCAS rocks from a deep continental source can be ruled out.

The scattering data points of dioritic members of both SUPS and KCAS in figures 3, 6, 8 and 9 suggests a common or at least similar source composition for both series, more enriched in HREE and depleted in Nb, Sr, P and Ti for the KCAS. The hypothetical source is characterized by high LILE/LREE and LREE/HFSE (high field strength elements) ratios, as well as by high Th/U and low Rb/Sr ratios. It is more enriched in incompatible elements compared to ocean island or island arc volcanics (MORRISON, 1980; THOMPSON and FOWLER, 1986; THOMPSON et al., 1984).

There are three possible explanations for the LILE-enrichment: (A) a mantle source enriched in Ba, Rb, Th and K, (B) contamination of mantle melts by partial melts of a subducted slab and (C) contamination by crustal melts. The model of an enriched subcontinental mantle source is widely used to explain trace element patterns of subduction-related potassic to ultrapotassic rocks (FOLEY et al., 1987; STILLE et al., 1989; THOMPSON and FOWLER, 1986; WHELLER et al., 1987). Such a mantle source would have elevated Rb/Sr, Th/U, La/Nb ratios and LILE concentrations. The trace element characteristics and the similarity of the REE patterns from diorite to granite would also argue for the hypothesis of an enriched mantle source for the SUPS. A large numbers of different models for upper mantle enrichment have been suggested so far. STILLE et al. (1989) derive Hercynian lamprophyres from a metasomatically enriched mantle, which is, however, depleted in K₂O and enriched in Nb. Other models involve the rise of volatiles or

melts into the upper mantle to produce magmas with primary hydrous mineral phases such as hornblende and mica (see FOLEY et al., 1987, for references). Such a model is consistent with the presented data set, because it allows the formation of high-K/Na ultrapotassic rocks by partial melting of a phlogopite-bearing peridotite. The high LILE concentrations as well as the elevated Ba/Sr ratios are the result of this enrichment process and therefore characteristic of the involved mantle source. The mantle enrichment by either volatiles or melts happened in connection with a subduction process of unknown age; it is a distinct process, separated by space and/or time, from contamination during the Hercynian subduction.

Partial melts of a subducted slab that contains pelagic sediments and oceanic crust can be highly enriched in incompatible elements (SCHREYER et al., 1987). Contamination with this material would lead to high Ba/La and Ba/Sr (STILLE et al., 1989). Ba/Sr ratios in the SUPS range from 2.4 up to 9.7, the crustal mean being around 0.9–1.1. Initial Sr ratios of the Givv syenite range between 0.7073 and 0.7076 (unpublished data) and indicate either a volumetrically subordinate crustal contamination or contamination by melts from a low-Rb/Sr source. Considerable contamination by crustal melts (C) would cause much higher initial Sr ratios and low concentrations of Sr, Eu, Ti and P. The lack of a negative Eu anomaly as well as La_N/Yb_N ratios that are typical for alkali basalts, argue against this possibility. Crustal contamination is also indicated for the SUPS by low initial Hf isotopic compositions (SCHALTEGGER and CORFU, in prep.).

The rocks of the KCAS do not show the typical LILE-enrichment. The elemental variation in figure 4 may be interpreted as contamination of high-Cr, Ni, Ca, Sr mantle melts by crustal melts rich in Zr and LILE and low in Ba. The hybrid melts experienced fractional crystallization of titanite, ilmenite, zircon, apatite, plagioclase, pyroxenes and hornblende that caused scatter of Ti, Zr, Ca, P, Sr, Cr, Ni. The observed element scatter may consequently be the result of the combination of both processes. A mixing process is indicated in figure 11 (Ba/Sr vs Zr/Th diagram): a common source with Ba/Sr around 2.5 and a Zr/Th of 2 is mixed with crustal melts, causing a steep increase of the Zr/Th ratio and lowering the Ba/Sr ratio, and on the other hand with partial melts from the subducted slab, producing high Ba/Sr ratios (STILLE et al., 1989) and not changing the Zr/Th ratio. An alternative interpretation would be fractional crystallization: Assuming a bulk distribution coefficient $D_{Ba/Sr} > 1$, the Ba/Sr ratio might be enhanced by increasing modal content of K-feldspar, indicated by the data points of the Central Aar granite. The Zr/Th ratio




Fig. 11 Zr/Th vs Ba/Sr diagram. Same symbols as in figure 2. An original mantle source with Ba/Sr = 3 and Zr/Th = 4 was enriched during a subduction process involving sediments and leading to elevated Ba/Sr ratios of the shoshonitic-ultrapotassic rocks. The latter may be increased by K-feldspar accumulation. The high-K calc-alkaline rocks were possibly contaminated by crustal melts and experienced allanite fractionation, increasing the Zr/Th ratio in the melt.

on the other hand increases with the removal of allanite from the melt; such a process is indicated by highly variable La/Nb ratios in the KCAS (see Fig. 4). Figure 11 leads to the conclusion that different processes or mixing endmembers are responsible for the development of each of the two investigated rock series.

Tectonic implications

The results of the geochemical investigations indicate that the dioritic, syenitic, monzonitic and granitic rocks of the eastern Aar massif are derived from an enriched subcontinental mantle. The geochemical composition, the contamination by crustal melts or fluids and the occurrence in an orogenic setting certainly argues for a subduction-related genesis at some stage of the Laurasia-Gondwana collision. It is generally accepted that ultrapotassic magmatism may either be generated above active Andean-type subduction zones or as well in collisional belts, shortly after the closure of the ocean basin (THOMPSON et al., 1984; THOMPSON and FOWLER, 1986).

The nature and polarity of Hercynian subduction is still controversial, but most of the current models imply the presence of subducted oceanic crust (STILLE et al., 1989; MERCOLLI and OBERHÄNSLI, 1988). No preserved Paleozoic oceanic crust has been detected in the central Alpine segment of the Hercynides up to now, thus the complete consumption of the ocean basin must be envisaged. The obvious lack of a Hercynian high-

grade high-T metamorphism or contemporaneous S-type magmatism favours the model of an Andean-type subduction zone rather than a classical continent-continent collision (FINGER and STEYRER, 1990). The active continental margin was formed by a magmatic arc that underwent considerable crustal shortening by upper crustal thin-skinned thrusting (OBERHÄNSLI et al., 1988) and possibly mid-crustal detachment faulting, as demonstrated by SHEFFELS (1990) for the Andes. The shoshonitic-ultrapotassic rocks are thought to be intruded into the thickened crust during a period of crustal relaxation or delamination.

The late Hercynian tectonic regime, creating strike-slip faults and pull-apart basins (BONIN, 1988; ZIEGLER, 1986), tapped the late-orogenic deep crustal magma chambers that generated large late Hercynian calc-alkaline batholiths in the Aar, Gotthard and Mont-Blanc massifs. The active continental margin had a more southern position at that time and the Andean-type subduction generated calc-alkaline magmas as young as Permian, e.g. the 262 Ma-old Lugano porphyries (STILLE and BULETTI, 1987). It is concluded that the onset of the slightly distensional, late Hercynian wrench tectonics is younger in the central region of the Hercynian fold belt than previously assumed, marked by the intrusion of the 334 Ma-old shoshonitic-ultrapotassic series of the Eastern Aar Massif.

Conclusions

Geochemical investigations on late Hercynian intrusive rocks of the Aar Massif lead to the following conclusions:

1. In the eastern Aar Massif, two different intrusive series exist that can be distinguished on geochemical basis: A shoshonitic-ultrapotassic suite, consisting of shoshonitic diorites, syenites and granites, is derived from an enriched mantle source; a high-K calc-alkaline series, consisting of shoshonitic diorites, monzodiorites and granites, is derived from the same or a similar source, but experienced extensive fractional crystallization and possibly contamination by crustal melts.

2. The subcontinental lithospheric mantle, which has been probed by the shoshonitic-ultrapotassic rocks, was geochemically enriched during a pre-Hercynian subduction event that created the characteristic geochemical features as LILE-enrichment and elevated Ba/Sr ratios, the latter due to the involvement of pelagic sediments.

3. The ultrapotassic rocks are typical of group III rocks of FOLEY et al. (1987) and have to be placed into an orogenic setting. Their generation

and emplacement is in direct relationship with an Upper Carboniferous Andean-type subduction.

4. The shoshonitic-ultrapotassic suite marks the beginning of late Hercynian strike-slip tectonics that created the pathways for the ascent of the voluminous late Hercynian magmatic rocks in the Aar Massif as well as in other parts of the Helvetic basement.

Acknowledgements

We wish to thank the staff of the Department of Mineralogy and Petrography in Bern, where the authors carried out their research, and of the departments in Fribourg and Lausanne, where parts of the analyses were performed. The paper was written during a fellowship of the Schweizerischer Nationalfonds to U.S. at the Royal Ontario Museum (Toronto, Canada), which is kindly acknowledged. The staff members of ROM Geochronology Lab are thanked for their help and hospitality. The manuscript benefitted of comments and fruitful discussions with F. Corfu and L. Heaman, both Toronto. The paper was reviewed by R. Oberhänsli, Mainz.

References

- BÖHM, C. (1988): Vulkanoklastite im östlichen Aarmassiv (Val Russein, Graubünden). *Schweiz. Mineral. Petrogr. Mitt.* 68, 501–508.
- BONIN, B. (1988): From orogenic to anorogenic environments: evidence from associated magmatic episodes. *Schweiz. Mineral. Petrogr. Mitt.* 68, 301–312.
- CHAUVEL, C. and JAHN, B.-M. (1984): Nd–Sr isotope and REE geochemistry of alkali basalts from the Massif Central, France. *Geochim. Cosmochim. Acta* 48, 93–110.
- CORRIVEAU, L. (1990): Proterozoic subduction and terrane amalgamation in the southwestern Grenville Province, Canada: Evidence from ultrapotassic to shoshonitic plutonism. *Geology* 15, 614–617.
- DEMPSTER, T. (1986): Isotope systematics in minerals: biotite rejuvenation and exchange during Alpine metamorphism. *Earth Planet. Sci. Lett.* 78, 355–367.
- ELLAM, R.M., HAWKESWORTH, C.J., MENZIES, M.A. and ROGERS, N.W. (1989): The volcanism of Southern Italy: Role of subduction and the relationship between potassic and sodic alkaline magmatism. *J. Geophys. Res.* 94, B4, 4589–4601.
- FINGER, F. and STEYRER, H.P. (1990): I-type granitoids as indicators of a late Paleozoic convergent ocean-continent margin along the southern flank of the central European Variscan orogen. *Geology* 18, 1207–1210.
- FOLEY, S.F., VENTURELLI, G., GREEN, D.H. and TOSCANI, L. (1987): The ultrapotassic rocks: Characteristics, classification and constraints for petrogenetic models. *Earth-Sci. Rev.* 24, 81–134.
- FRANKS, G.D. (1968a): A study of upper Paleozoic sediments and volcanics in the northern part of the eastern Aar Massif. *Eclogae geol. Helv.* 61, 49–140.
- FRANKS, G.D. (1968b): The pre-Westphalian (Hercynian) metamorphism and structures of the Tödi area (Aar Massif). *Schweiz. Mineral. Petrogr. Mitt.* 48, 667–694.
- FREY, M., TEICHMÜLLER, M., TEICHMÜLLER, R., MULLIS, J., KÜNZI, B., BREITSCHMID, A., GRUNER, U. and SCHWIZER, B. (1980): Very low-grade metamorphism in external parts of the Central Alps: Illite crystallinity, coal rank and fluid inclusion data. *Eclogae geol. Helv.* 73, 173–203.
- FRISCH, W., MÉNOT, R.-P., NEUBAUER, F. and VON RAUMER, J.F. (1990): Correlation and evolution of the Alpine basement. *Schweiz. Mineral. Petrogr. Mitt.* 70, 265–286.
- GNOS, E. (1988): *Geologie und Petrographie des westlichen Brunnitals (Maderanertal, Kanton Uri)*. Unpubl. diploma thesis, Univ. of Bern, 97 pp.
- HUBER, W. (1948): *Petrographisch-mineralogische Untersuchungen im südöstlichen Aarmassiv*. *Schweiz. Mineral. Petrogr. Mitt.* 28, 555–642.
- KÜPFER, Th. (1977): *Mineralogisch-petrographische und geochemische Untersuchung der Syenodiorite und der Ganggesteine im Puntegliasgebiet (GR)*. Unpubl. Ph. D. thesis, Univ. of Bern.
- LABHART, T.P. (1977): *Aarmassiv und Gotthardmassiv*. *Sammlg. Geol. Führer* 63, Bortraeger Berlin-Stuttgart, 173 pp.
- LABHART, T.P. and RYBACH, L. (1971): Abundance and distribution of uranium and thorium in the syenite of Piz Giuf (Aar Massif, Switzerland). *Chem. Geol.* 7, 237–251.
- MARQUER, D. (1987): *Transfert de matière et déformation progressive des granitoïdes. Exemple des massifs de l'Aar et du Gothard (Alpes Centrales suisses)*. *Mém. Docum. Centre Arm. Ét. Struct. Socles, Rennes*, 10, 287 pp.
- MERCOLLI, I. and OBERHÄNSLI, R. (1988): Variscan tectonic evolution in the Central Alps: a working hypothesis. *Schweiz. Mineral. Petrogr. Mitt.* 68, 491–500.
- MITTFELHELDT, D.W. and MILLER, C.F. (1983): Geochemistry of the Sweetwater Wash Pluton, California: Implications for "anomalous" trace element behaviour during differentiation of felsic magmas. *Geochim. Cosmochim. Acta* 47, 109–124.
- MORRISON, G.W. (1980): Characteristics and tectonic setting of the shoshonite rock association. *Lithos*, 13, 97–108.
- NAKAMURA, N. (1974): Determination of REE, Ba, Mg, Na and K in carbonaceous and ordinary chondrites. *Geochim. Cosmochim. Acta* 38, 757–775.
- OBERHÄNSLI, R. (1986): Geochemistry of meta-lamprophyres from the Central Swiss Alps. *Schweiz. Mineral. Petrogr. Mitt.* 66, 315–342.
- OBERHÄNSLI, R., SCHENKER, F. and MERCOLLI, I. (1988): Indications of Variscan nappe tectonics in the Aar Massif. *Schweiz. Mineral. Petrogr. Mitt.* 68, 509–520.
- OBERHÄNSLI, R., KRÄHENBÜHL, U. and STILLE, P. (1991): Contrasting REE characteristics in meta-lamprophyres from Variscan Massifs of the Central Swiss Alps and REE patterns in lamprophyres from Variscan terranes of Western Europe. *Schweiz. Mineral. Petrogr. Mitt.* 71, 53–62.
- PECERILLO, A. and TAYLOR, S.R. (1976): Geochemistry of Eocene calc-alkaline volcanic rocks from the Kastamonu area, Northern Turkey. *Contrib. Mineral. Petrogr.* 58, 63–81.
- SCHALTEGGER, U. (1989): *Geochemische und isotopengeologische Untersuchungen am Zentralen Aaregranit und seinen assoziierten Gesteinen zwischen Aare und Reuss (Aarmassiv, Schweiz)*. Unpubl. Ph. D. thesis University of Bern, 144 pp.

- SCHALTEGGER, U. (1990): The Central Aar Granite: Highly differentiated calc-alkaline magmatism in the Aar Massif (Central Alps, Switzerland). *Eur. J. Mineral.* 2, 254–259.
- SCHALTEGGER, U. and CORFU, F.: An enriched subcontinental mantle source for late Hercynian magmatism in the Central Alps: Evidence from precise U–Pb ages and initial Hf isotopes (manuscript in preparation).
- SCHALTEGGER, U. and KRÄHENBÜHL, U. (1990): Heavy rare earth enrichment in granites of the Aar Massif (Central Alps, Switzerland). *Chem. Geol.* 89, 1/2, 49–63.
- SCHALTEGGER, U. and VON QUADT, A. (1990): U–Pb zircon dating of the Central Aar Granite (Aar Massif, Switzerland). *Schweiz. Mineral. Petrogr. Mitt.* 70, 361–371.
- SCHALTEGGER, U., CORFU, F. and KROGH, T.E. (1991): Precise U–Pb ages and initial Hf isotopic compositions of late Hercynian mafic to acid intrusives in the Aar Massif, Central Alps. *Terra abstracts* 3, 37.
- SCHENKER, F. (1986): Spätpaläozoischer saurer Vulkanismus und Beckenbildung im Aarmassiv unter kompressiver Tektonik. Unpubl. Ph. D. thesis, Univ. of Bern. 116 pp.
- SCHREYER, W., MASSONNE, H.J. and CHOPIN, C. (1987): Continental crust subducted to depths near 100 km: implications for magma and fluid genesis in collision zones. In: MYSEN, B.O. (ed.), *Magmatic Processes: Physicochemical Principles*. *Geochem. Soc. Spec. Publ.* 1, 155.
- SEEMANN, U. (1975): Mineralogisch-petrographische und geochemische Untersuchungen der granitischen Gesteine der Val Punteglias (GR). *Schweiz. Mineral. Petrogr. Mitt.* 55, 257–306.
- SHEFFELS, B.M. (1990): Lower bound on the amount of crustal shortening in the central Bolivian Andes. *Geology* 18, 812–815.
- SHIREY, S.B. and HANSON, G.N. (1984): Mantle-derived Archean monzodiorites and trachyandesites. *Nature*, Nr. 310, 222–224, July 19, 1984.
- SIGRIST, F. (1947): Beitrag zur Kenntnis der Petrographie und der alpinen Zerrklüftlagerstätten des östlichen Aarmassivs. *Schweiz. Mineral. Petrogr. Mitt.* 27, 39–183.
- STILLE, P. (1988): *Geochemische Aspekte der Krustenentwicklung im zentral- und südalpinen Raum*. Habilitationsschrift ETH Zürich.
- STILLE, P. and BULETTI, M. (1987): Nd–Sr isotopic characteristics of the Lugano volcanic rocks and constraints on the continental crust formation in the South Alpine domain (N-Italy-Switzerland). *Contrib. Mineral. Petrol.* 96, 140–150.
- STILLE, P., OBERHÄNSLI, R. and WENGER-SCHENK, K. (1989): Hf–Nd and trace element constraints on the genesis of alkaline and calc-alkaline lamprophyres. *Earth Planet. Sci. Lett.* 96, 209–219.
- SUN, S.-S. (1980): Lead isotope study of young volcanic rocks from mid-ocean ridges, ocean islands and island arcs. *Phil. Trans. R. Soc. London, A* 297, 409–445.
- THOMPSON, R.N. (1982): Magmatism of the British Tertiary province. *Scottish J. Geol.* 18, 49–107.
- THOMPSON, R.N. and FOWLER, M.B. (1986): Subduction-related shoshonitic and ultrapotassic magmatism: a study of Siluro-Ordovician syenites from the Scottish Caledonides. *Contrib. Mineral. Petrol.* 94, 507–522.
- THOMPSON, R.N., MORRISON, M.A., HENDRY, G.L. and PARRY, S.J. (1984): An assessment of the relative roles of crust and mantle in magma genesis: an elemental approach. *Phil. Trans. R. Soc. London, A* 310, 549–590.
- THOMPSON, R.N., LEAT, P.T., DICKIN, A.T., MORRISON, M.A., HENDRY, G.L. and GIBSON, S.A. (1989): Strongly potassic mafic magmas from lithospheric mantle sources during continental extension and heating: evidence from Miocene minettes of north-west Colorado, U.S.A. *Earth Planet. Sci. Lett.* 98, 139–153.
- WHELLER, G.E., VARNE, R., FODEN, J.D. and ABBOTT, M.J. (1987): Geochemistry of quaternary volcanism in the Sunda-Banda Arc, Indonesia, and three-component genesis of island-arc basaltic magmas. *J. Volcanol. Geotherm. Res.* 32, 137–160.
- ZIEGLER, P.A. (1986): Geodynamic model for the Paleozoic crustal consolidation of Western and Central Europe. *Tectonophysics* 126, 303–328.

Manuscript received March 1, 1991; revised manuscript accepted July 15, 1991.