

Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =
Bulletin suisse de minéralogie et pétrographie

Band: 72 (1992)

Heft: 3

Rubrik: New minerals recently approved by the Commission on New Minerals
and Mineral Names, International Mineralogical Association

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 13.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

**New minerals recently approved
by the
Commission on New Minerals and Mineral Names
International Mineralogical Association**

The information given here is provided by the Commission on New Minerals and Mineral Names, I.M.A. for comparative purposes and as a service to mineralogists working on new species.

Each mineral is described in the following format:

IMA No.

(any relationship to other minerals)

Chemical Formula

Crystal system, space group

unit cell parameters

Colour; lustre; diaphaneity.

Optical properties

Strongest lines in the X-ray powder diffraction pattern

The names of these approved species are considered confidential information until the authors have published their descriptions or released information themselves.

No other information will be released by the commission

J. A. Mandarino, Chairman

Commission on New Minerals and Mineral Names

International Mineralogical Association

1991 Proposals

IMA No. 91-001

$\text{Hg}_2^{+1}\text{Hg}_3^{+2}\text{Cr}^{+6}\text{O}_5\text{S}_2^{-2}$

Triclinic: $P\bar{1}$

a 8.116, b 9.501, c 6.891 Å, α 100.43°,

β 110.24°, γ 82.80°

Orange-red; adamantine; transparent.

Biaxial (sign unknown), all indices of refraction are greater than 2.

5.72 (90), 3.373 (60), 3.008 (100), 2.864 (50b),
2.774 (50), 2.536 (50), 2.486 (50), 2.425 (60).

Biaxial (+), α 2.38, β 2.42, γ 2.47, 2V(calc.) 85°.

3.164 (100), 2.934 (90), 2.842 (45), 2.495 (45),
1.769 (45), 1.734 (80).

IMA No. 91-005

$(\text{Zn,Co,Ni})_6(\text{SO}_4)(\text{OH,Cl})_{10} \cdot 5 \text{H}_2\text{O}$

Hexagonal: $P6_3$, $P6_3/m$ or $P6_322$

a 8.344, c 21.59 Å

Bright to deep pink; vitreous to pearly;
transparent.

Uniaxial (-), ω 1.584, ϵ 1.544

10.8 (100), 3.300 (90), 2.725 (60), 2.563 (50),
2.351 (40), 1.575 (30).

IMA No. 91-003

The niobium analogue of bismutotantalite.

$\text{Bi}(\text{Nb,Ta})\text{O}_4$

Orthorhombic: $Pcmm$

a 4.992, b 5.677, c 11.731 Å

Black; semi-metallic; transparent in small
(< 0.03 mm) fragments.

IMA No. 91-007

$\text{Mn}_3(\text{OH})_4(\text{VO}_4)_2$


Monoclinic: $C2/m$

a 9.604, b 9.558, c 5.393 Å, β 98.45°

Orange-red; vitreous; transparent.
Biaxial (-), α' 1.803, γ' 1.810, $2V(\text{meas.})$ large.
4.76 (S), 3.00 (M), 2.680 (VS), 2.656 (M),
2.155 (M), 1.565 (M), 1.510 (M).

IMA No. 91-008

The Ba-dominant end-member of the alunite group.



Hexagonal: $R\bar{3}m$

a 6.992, c 17.22 Å


White to light yellowish; vitreous; transparent.

Uniaxial (+), ω 1.588, ε 1.604.

5.73 (50), 3.49 (55), 2.98 (100), 2.283 (80),
1.909 (70), 1.747 (60).

IMA No. 91-009

The Ca-dominant end-member of the alunite group.



Hexagonal: $R\bar{3}m$


a 6.983, c 16.759 Å

White to light yellowish; vitreous; transparent.

Uniaxial (+), indices of refraction unknown.

4.91 (69), 2.97 (100), 2.231 (51), 1.899 (43),
1.745 (37), 1.375 (40).

IMA No. 91-010



Hexagonal: $R\bar{3}m$


a 6.240, c 26.784 Å

Yellow-brown; vitreous; transparent.

Uniaxial (-), ω 1.770, ε 1.759.

5.00 (60), 3.119 (100), 2.689 (80), 2.558 (100),
2.505 (80), 1.560 (80).

IMA No. 91-012



Monoclinic: $P2_1/m$

a 28.77, b 7.01, c 13.72(2) Å, β 96.6(2)°

Orange-yellow; vitreous; transparent.

Biaxial (-), α 1.696, β 1.711, γ 1.715,

$2V(\text{meas.})$ 57°, $2V(\text{calc.})$ 54°.

3.41 (8), 3.22 (8), 2.83 (10), 2.81 (10), 2.24 (7),
1.750 (6).

IMA No. 91-013



Orthorhombic: $Cmcm$, $Cmc2_1$ or $C2cm$

a 29.77, b 11.03, c 17.111(5) Å


Colourless (white or grey in aggregates);
vitreous; transparent.

Biaxial (-), α 1.532, β 1.548, γ 1.559(2),

$2V(\text{meas.})$ 79°, $2V(\text{calc.})$ 79°.

10.38 (100), 4.516 (75), 3.220 (65), 3.097 (80),
2.972 (65), 2.773 (90).

IMA No. 91-014



Triclinic: $P\bar{1}$

a 10.244, b 11.924, c 5.276 Å, α 103.491°,
 β 96.960°, γ 91.945°


Olive-green with brownish or yellowish shades;
vitreous; transparent.

Biaxial (+), α 1.569, β 1.574, γ 1.590,

$2V(\text{meas.})$ 58°, $2V(\text{calc.})$ 59°.

11.57 (100), 3.386 (19), 3.006 (21), 2.992 (28),
2.716 (22), 2.598 (26).

IMA No. 91-015



Monoclinic: $P2_1/c$

a 24.91, b 11.94, c 14.92 Å, β 94.47(9)°

Colourless; vitreous; transparent.


Biaxial (-), α 1.460, β 1.478, γ 1.481,

$2V(\text{meas.})$ 43°, $2V(\text{calc.})$ 44°.

4.26 (60), 3.08 (100), 2.938 (70B), 2.649 (60B),
2.400 (35), 2.289 (35).

IMA No. 91-016

A member of the adelite-descloizite group.



Orthorhombic: $P2_12_12_1$

a 5.838, b 7.224, c 8.690(1) Å


Deep red; vitreous; transparent.

Biaxial (+), α 1.840, β (calc.) 1.854, γ 1.920,
 $2V(\text{meas.})$ 50°.

5.558 (S), 3.070 (S), 2.687 (S), 2.584 (VS),
1.565 (M).

IMA No. 91-017

The ferric-analogue of philipsbornite.



Hexagonal: $R\bar{3}m$

a 7.359, c 17.113(8) Å


Greenish-yellow; vitreous to adamantine;
translucent to transparent.

Uniaxial (-), ω 1.975, ε 1.955.

5.966 (50), 3.678 (40), 3.092 (100), 2.283 (30),
1.992 (30), 1.840 (25).

IMA No. 91-018

The Mg-dominant analogue of congolite and
the rhombohedral dimorph of boracite.



Hexagonal: $R3c$


a 8.574, c 20.99 Å

Colourless; vitreous; transparent.

Uniaxial (-), ω 1.684, ε 1.668.


3.497 (34), 3.028 (100), 2.711 (66), 2.144 (37),
2.050 (73), 1.828 (25).

IMA No. 91-019



- Monoclinic: $P2_1/m$
 a 10.625, b 3.264, c 8.990 Å, β 109.15°.
 Dark brown to black; submetallic to metallic;
 opaque (translucent in thin fragments).
 In reflected light: creamy white (in oil, white
 with a weak brown tint); no internal
 reflections; anisotropy visible along grain
 boundaries (in oil, clearly visible); bire-
 flectance not visible (in oil, very weak
 along grain boundaries); nonpleochroic.
 R-values: (15.5–15.9%) 470 nm, (15.0–
 15.5%) 546 nm, (14.8–15.0%) 589 nm,
 (14.2–14.5%) 650 nm.
 2.985 (67), 2.811 (94), 2.749 (100), 2.391 (85),
 1.779 (48), 1.709 (35).
- IMA No. 91-020
 $Ba(Al,Mg)(PO_4,CO_3)(OH)_2 \cdot H_2O$
 Orthorhombic: Pnna or Pnnc
 a 8.939, b 5.669, c 11.073(3) Å
 Pale blue; silky; translucent.
 Biaxial (-), α 1.616, β 1.629, γ 1.640,
 2V(meas.) 70–90°, 2V(calc.) 85°.
 5.54 (79), 3.479 (82), 3.345 (59), 2.768 (100),
 2.543 (61), 2.072 (41).
- IMA No. 91-021
 A polymorph of mundrabiliaite.
 $Ca(NH_4)_2(HPO_4)_2 \cdot H_2O$
 Orthorhombic: space group unknown
 a 20.959, b 7.403, c 6.478(5) Å
 White; vitreous; transparent.
 Biaxial (-), α 1.506, β 1.510, γ 1.512,
 2V(meas.) 65°, 2V(calc.) 70°.
 10.5 (57), 6.99 (100), 4.739 (36), 3.705 (89),
 3.651 (39), 3.177 (55).
- IMA No. 91-022
 $KZr_2(PO_4)_3$
 Hexagonal: $R\bar{3}c$
 a 8.687, c 23.877(7) Å
 Pale blue to blue-green to nearly colourless;
 vitreous; transparent.
 Uniaxial (+), ω 1.656, ϵ 1.682.
 6.41 (50), 4.679 (50), 4.329 (100), 3.806 (90),
 2.928 (90), 2.502 (50).
- IMA No. 91-023
 The Cs-analogue of gainesite.
 $NaCs(Be,Li)Zr_2(PO_4)_4$
 Tetragonal: $I4_1/amd$
 a 6.573, c 17.28 Å
 White to colourless; vitreous; translucent to
 transparent.
 Uniaxial (+), ω 1.634, ϵ 1.645.
 6.159 (90), 4.326 (80), 4.099 (40), 3.281 (80),
 3.060 (100), 2.896 (30), 1.849 (30).
- IMA No. 91-024
 $Sb_6(SO_4)O_8 \cdot H_2O$
 Triclinic: P1
 a 11.434, b 29.77, c 11.314(4) Å, α 91.07°,
 β 119.24°, γ 92.82°
 Colourless to white; adamantine; transparent
 to translucent.
 Biaxial (+), mean n 2.08, birefringence low,
 2V(meas.) \gg 60°.
 14.835 (50), 9.270 (41), 6.810 (67), 3.304 (93),
 3.200 (39), 3.092 (100).
- IMA No. 91-025
 $Cu_4(As,Sb)_2S_5$
 Orthorhombic: space group unknown
 a 14.51, b 13.30, c 17.96(1) Å
 Silvery lead grey; metallic; opaque.
 In reflected light: grey, weak anisotropism,
 weak bireflectance, nonpleochroic. R_{min}
 and R_{max} : (31.5, 32.5%) 470 nm, (31.1,
 32.0%) 546 nm, (30.3, 31.15%) 589 nm,
 (27.2, 23.4%) 650 nm.
 3.36 (7), 2.999 (100), 2.594 (20), 2.238 (6),
 1.833 (40), 1.564 (15b).
- IMA No. 91-026
 $(Co,Zn,Ni)_6(SO_4)(OH,Cl)_{10} \cdot 8 H_2O$
 Hexagonal: space group unknown
 a 8.363, c 26.18(7) Å
 Pink to light pink; pearly; transparent.
 Uniaxial (-), ω 1.568, ϵ 1.542.
 13.1 (100), 3.523 (30), 2.985 (30), 2.681 (40),
 2.527 (90).
- IMA No. 91-027
 The fluorine-analogue of britholite-(Ce) of the
 apatite group.
 $(REE, Ca)_5(Si,P)_3O_{12}F$, where Ce is the
 dominant REE.
 Hexagonal: $P6_3/m$
 a 9.517, c 6.983(4) Å
 Tan, reddish-brown; adamantine; opaque to
 translucent.
 Uniaxial (-), ω 1.792, ϵ 1.786.
 2.845 (100), 2.822 (40), 2.747 (30), 1.970 (30),
 1.870 (40).
- IMA No. 91-028
 A member of the amphibole group.
 $(Na,K)Na_2[Mg_2(Fe^{3+},Mn^{3+})_2Li]Si_8O_{22}(OH,F)_2$
 Monoclinic: $C2/m$
 a 9.808, b 17.850, c 5.289(1) Å, β 104.22(2)°
 Dark red; vitreous; translucent.
 Biaxial (+), α 1.667, β 1.675, γ 1.691,
 2V(meas.) 59–71°, 2V(calc.) 71°.
 8.399 (56), 3.383 (18), 3.254 (20), 3.122 (100),
 2.798 (48), 2.696 (15).

IMA No. 91-029


Hexagonal: $\text{P}\bar{3}1\text{m}$, $\text{P}31\text{m}$ or $\text{P}312$

a 16.016, c 9.789(2) Å

Light-blue; vitreous; transparent.

Uniaxial (+), ω 1.600, ϵ 1.605.4.6195 (100), 3.3537 (100), 2.3431 (80),
2.0909 (60), 1.8050 (70), 1.7496 (60).

IMA No. 91-030


Tetragonal: $\text{I}4/\text{mmm}$, $\text{I}42\text{m}$, $\text{I}4\text{m}2$, $\text{I}4\text{mm}$
or $\text{I}422$

a 3.9922, c 22.514(2) Å

Carmine; adamantine; translucent.

In reflected light: grey, weak to moderate
anisotropy, moderate bireflectance, weak
pleochroism, internal reflections abundant,
 R_1 and R_2 (19.6, 22.0%) 470 nm, (18.0,
20.5%) 546 nm, (17.4, 19.6%) 589 nm,
(16.95, 18.8%) 650 nm.3.507 (32), 2.983 (100), 2.816 (78), 1.989 (75),
1.658 (51), 1.586 (33).

IMA No. 91-031

Monoclinic: $\text{P}2_1/\text{c}$ (pseudo $\text{C}2/\text{c}$)a 15.122, b 14.358, c 22.063 Å, β 108.68°.


Dark blue; vitreous; transparent.

Biaxial (-), α 1.590, β 1.610, γ 1.619,

2V(meas.) 65°, 2V(calc.) 67°.

3.393 (100), 3.368 (55), 3.200 (53), 3.188 (65),
3.120 (85), 3.098 (57).

IMA No. 91-032

Orthorhombic Immm (pseudocubic)

a 7.544, b 7.558, c 7.560(4) Å

Dark bottle green; vitreous to adamantine;
transparent.Biaxial (-), the indices of refraction are
between 1.92 and 1.94.3.774 (100), 2.671 (35), 2.395 (30), 1.904 (15),
1.697 (60), 1.548 (40).


IMA No. 91-033

Cubic: $\text{Fd}3\text{m}$

a 7.933(5) Å

Colour unknown because of the small grain
size; metallic; opaque.In reflected light: silvery grey, dark grey when
highly oxidized; no anisotropy, bireflectance,
pleochroism or internal reflections;
 R (56.0%) 470 nm, (59.5%) 546 nm,
(60.0%) 589 nm, (62.0%) 650 nm.4.595 (21), 2.810 (30), 2.391 (100), 2.301 (25),
1.526 (23), 1.196 (26).

IMA No. 91-034

Orthorhombic: Pmmn , $\text{Pmn}2_1$ or $\text{P}2_1\text{nm}$

a 15.337, b 17.051, c 6.931 Å

Canary yellow; vitreous; transparent.

Biaxial (-), α 1.603(calc.), β 1.690, γ 1.710,
2V(meas.) 49°.8.55 (100), 6.94 (50), 4.11 (60), 3.723 (60),
3.460 (50), 2.772 (70).

IMA No. 91-037

Monoclinic: $\text{C}2/\text{m}$ or Cm a 13.515, b 4.098, c 26.000 Å, β 93.00°

Grey; metallic; opaque.

In reflected light: white, distinct anisotropy,
very weak bireflectance, no pleochroism,
no internal reflections, R_{max} and R_{min} .
(42.2, 39.7%) 470 nm, (41.4, 38.8%) 546 nm,
(40.8, 37.9%) 589 nm, (39.8, 36.9%) 650 nm.3.49 (8), 3.37 (9), 3.24 (9), 2.82 (10), 2.01 (7),
1.992 (8), 1.967 (6).

IMA No. 91-038


Hexagonal: $\text{P}6_3/\text{mmc}$, $\text{P}6_3\text{mc}$ or $\text{P}62\text{c}$

a 5.951, c 33.358 Å

Black; submetallic; opaque.

In reflected light: grey with pale brownish tint,
moderate anisotropy, weak bireflectance,
no pleochroism, no internal reflections, R_{O}
and R_{E} . (23.6, 22.3%) 470 nm, (22.8, 21.9%)
546 nm, (22.2, 21.5%) 589 nm, (21.3, 21.0%)
650 nm.4.168 (55), 3.011 (60), 2.9750 (70), 2.8017 (95),
2.6236 (100), 2.6125 (90).


IMA No. 91-042

Orthorhombic: $\text{I}222$

a 8.984, b 8.937, c 8.927 Å

White, colourless, light yellow; vitreous;
transparent.Biaxial (-), α 1.529, β (calc.) 1.530, γ 1.531,
2V(meas.) 76°.6.33 (8), 4.46 (8), 3.66 (10), 2.60 (8), 1.760 (8),
1.351 (8).

IMA No. 91-043

The Sb-dominant member of the colusite
group.Cubic: $\text{P}43\text{n}$

a 10.705 Å

Colour not observed because of the small size;
metallic; opaque.

- In reflected light: grey with a light-brown tint; no anisotropy, bireflectance, pleochroism or internal reflections; R (25.2%) 470 nm, (28.3%) 546 nm, (29.9%) 589 nm, (31.0%) 650 nm.
3.10 (10), 1.892 (9), 1.614 (7), 1.226 (4), 1.094 (6), 1.030 (4).
- IMA No. 91-044
The Ge-dominant member of the colusite group.
 $\text{Cu}_{26}\text{V}_2(\text{Ge,As})_6\text{S}_{32}$
Cubic: P43n
a 10.568 Å
Grey-black; metallic; opaque.
In reflected light: greenish-yellow, olive-yellowish-cream; no internal reflections, anisotropy, bireflectance or pleochroism; R (23.8%) 470 nm, (27.3%) 546 nm, (27.9%) 589 nm, (27.9%) 650 nm.
3.05 (10), 2.64 (4), 1.870 (5), 1.595 (3), 1.320 (3), 1.212 (3), 1.079 (3), 1.017 (5).
- IMA No. 91-045
 $(\text{Ca,Mg})_{14}\text{Mn}_3\text{Zn}_3\text{Be}_6\text{Si}_{14}\text{O}_{50}(\text{OH,F})_8$
Monoclinic: P2₁/c
a 9.08, b 18.03, c 14.59(4) Å, β 104.8°
Colourless; vitreous; transparent.
Biaxial (-), α 1.674, β 1.680, γ 1.681, 2V(meas.) 29.0°, 2V(calc.) 44°.
2.863 (100), 2.771 (40), 2.653 (50), 2.388 (50), 2.272 (30), 1.832 (30).
- IMA No. 91-046
The Cu-dominant analogue of geigerite and chudobaite.
 $(\text{Cu,Co})_5(\text{AsO}_4)_2(\text{AsO}_3\text{OH})_2 \cdot 10 \text{H}_2\text{O}$
Triclinic: P1 or P1
a 8.033, b 10.374, c 6.446(5) Å, α 79.62°, β 84.95°, γ 86.21°
Green; vitreous; transparent.
Biaxial (+), α 1.634, β 1.662, γ 1.720, 2V(meas.) 75°, 2V(calc.) 72°.
10.2 (100), 8.01 (60), 4.001 (50), 3.667 (60), 3.151 (50), 3.063 (50).
- IMA No. 91-047
 Ti_3As_4
Orthorhombic: Pnma
a 8.894, b 10.855, c 9.079 Å
Dark red; adamantine to submetallic; opaque to translucent.
In reflected light: red, red internal reflections, strong anisotropy, strong bireflectance, no pleochroism. R_{max.} and R_{min.} are: (4.78, 3.93%) 481 nm, (4.64, 3.86%) 547 nm, (8.64, 7.81%) 591 nm, (13.72, 11.78%) 644 nm.
- 4.14 (M), 3.99 (S), 3.80 (M), 3.47 (MSb), 3.35 (M), 2.813 (VS), 2.537 (M), 2.264 (MSb).
- IMA No. 91-048
 $\text{Na}_{25}\text{Ba}(\text{Y,Gd,Dy})_2(\text{CO}_3)_{11}(\text{HCO}_3)_4(\text{SO}_4)_2\text{F}_2\text{Cl}$
Hexagonal: P6₃/m
a 8.811, c 37.03(3) Å
Light green to yellowish-green; vitreous; transparent.
Uniaxial (-), ω 1.536, ε 1.510.
4.79 (42), 3.32 (40), 2.829 (100), 2.659 (51b), 2.531 (71b), 2.270 (90).
- IMA No. 91-050
 $(\text{Ca,REE,Th})_{15}\text{As}^{+5}(\text{As}_{0.5}^{+3}\text{Na}_{0.5})\text{Fe}^{+3}\text{Si}_6\text{B}_4\text{O}_{40}\text{F}_7$
Hexagonal: R3m
a 10.795, c 27.336(4) Å
Yellowish-green; vitreous; transparent.
Uniaxial (-), ω 1.757, ε 1.722.
2.993 (S), 2.950 (S), 1.839 (MS), 1.802 (MS), 1.686 (MS), 1.572 (MS).
- IMA No. 91-051
 $\text{Ag}_9\text{SbTe}_3(\text{S,Se})_3$
Monoclinic: P2, P2/m or Pm
a 8.900, b 8.302, c 19.49 Å, β 82.98°
Colour unknown because of the small grain size; metallic; opaque.
In reflected light: grey with faint green-blue hue, anisotropy present with brownish-grey tone, weak bireflectance, no pleochroism, no internal reflections, R_{max.} and R_{min.} (38.0, 34.2%) 470 nm, (36.6, 32.2%) 546 nm, (35.7, 31.8%) 589 nm, (34.0, 30.2%) 650 nm.
3.82 (6), 2.89 (4), 2.83 (4), 2.22 (10), 2.14 (3), 2.13 (4).
- IMA No. 91-052
The Sb-analogue of skutterudite.
 CoSb_3
Cubic: Im3
a 9.0411 Å
Tin-white; metallic; opaque.
In reflected light: tin-white, isotropic, no bireflectance, nonpleochroic, no internal reflections. R (59.0%) 470 nm, (58.7%) 546 nm, (58.7%) 589 nm, (58.7%) 650 nm.
2.85 (100), 2.01 (80), 1.92 (80), 1.84 (80), 1.50 (80), 1.185 (80), 1.147 (80), 0.780 (100).
- IMA No. 91-053
 $\text{Zn}_{12}(\text{CO}_3)_3(\text{SO}_4)(\text{OH})_{16}$
Orthorhombic: P22₁2
a 15.724, b 6.256, c 5.427(5) Å
White; vitreous; translucent.
Biaxial (probably +), α 1.635(3), β 1.650(3), γ could not be measured, 2V about 60°.

15.44 (100), 7.88 (100), 5.25 (20), 2.714 (40),
2.577 (20), 2.397 (20), 1.565 (30b).

IMA No. 91-054

$\text{Na}_{26}\text{Ce}_6(\text{SiO}_3)_6(\text{PO}_4)_6(\text{CO}_3)_6(\text{SO}_2)\text{O}$

Hexagonal: $\text{R}\bar{3}$

a 16.025, c 19.773 Å

Colourless to pale brown; vitreous; transparent.

Uniaxial (-), ω 1.589, ϵ 1.586.

8.076 (80), 6.544 (90), 4.659 (75), 3.776 (90),
3.159 (85), 2.683 (100).

IMA No. 91-055

A member of the epidote group, related to dollaseite-(Ce).

$(\text{Ca,REE})\text{REE}(\text{Mg,Fe})\text{MnAlSi}_3\text{O}_{11}(\text{OH})(\text{F,O})$

Monoclinic: $\text{P}2_1/\text{m}$

a 8.903, b 5.748, c 10.107 Å, β 113.41°

Dark greyish-brown; vitreous; transparent.

Biaxial (-), α 1.773, β 1.790, γ 1.803,

$2V(\text{meas.})$ 83°, $2V(\text{calc.})$ 82°.

9.32 (2), 5.23 (2), 4.67 (2), 3.52 (4), 2.91 (10),
2.73 (7), 2.63 (8), 1.437 (2).