

Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =
Bulletin suisse de minéralogie et pétrographie

Band: 74 (1994)

Heft: 2

Vereinsnachrichten: New minerals recently approved by the Commission on New
Minerals and Mineral Names International Mineralogical Association

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 13.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

**New minerals recently approved
by the
Commission on New Minerals and Mineral Names
International Mineralogical Association**

The information given here is provided by the Commission on New Minerals and Mineral Names, I.M.A. for comparative purposes and as a service to mineralogists working on new species.

Each mineral is described in the following format:

IMA No.

(any relationship to other minerals)

Chemical Formula

Crystal system, space group

unit cell parameters

Colour; lustre; diaphaneity

Optical properties

Strongest lines in the X-ray powder diffraction pattern

The names of these approved species are considered confidential information until the authors have published their descriptions or released information themselves.

No other information will be released by the commission.

J. A. Mandarino, Chairman
Commission on New Minerals and Mineral Names
International Mineralogical Association

1992 Proposals

IMA No. 92-001

$\text{FeZr}(\text{PO}_4)_2 \cdot 4 \text{H}_2\text{O}$

Monoclinic: $P2_1/c$

a 9.12, b 5.42, c 19.17 Å, β 94.8°

Pale yellowish white; vitreous to dull;
transparent.

Biaxial (+), α 1.644, β 1.652, γ 1.652,
 $2V(\text{meas.}) 0^\circ$, $2V(\text{calc.}) 0^\circ$.

9.58 (75), 4.572 (65), 4.382 (80), 4.092 (60),
3.160 (100), 2.640 (70).

Biaxial, indices of refraction calculated from
reflectance data at 589 nm: R_1 1.91, R_2 1.99.
3.644 (60), 3.466 (60), 3.206 (100), 2.924 (70),
2.782 (50), 1.984 (90).

IMA No. 92-003

The selenium analogue of stibnite.

Sb_2Se_3

Orthorhombic: $Pbnm$

a 11.593, b 11.747, c 3.984 Å

Black; metallic; opaque.

In reflected light: white, distinct anisotropism,
distinct bireflectance, pleochroic white to
greyish white. R_{max} & R_{min} : (42.62, 40.55%)
470 nm, (41.95, 39.02%) 546 nm, (41.23,
39.42%) 589 nm, (44.39, 41.56%) 650 nm.

IMA No. 92-002

$\text{Bi}_2\text{O}(\text{OH})_2\text{SO}_4$

Monoclinic: $P2_1/c$

a 7.700, b 13.839, c 5.686 Å, β 109.11°

Colourless; adamantine; transparent.

3.70 (70), 3.17 (50), 2.870 (100), 2.625 (60),
1.930 (30), 1.764 (35).

IMA No. 92-005

$\text{Mg}[\text{UO}_2(\text{AsO}_3)_x(\text{AsO}_4)_{1-x}]_2 \cdot 7 \text{H}_2\text{O}$, x about $2/3$
Monoclinic: $C2/m$

a 18.194, b 7.071, c 6.670 Å, β 99.70°

Bright yellow to straw yellow; vitreous;
transparent.

Biaxial (–), α 1.610, β 1.730, γ 1.740,
 $2V(\text{meas.})$ 34°, $2V(\text{calc.})$ 30°.

9.02 (100), 4.90 (40), 4.48 (80), 4.00 (40),
3.53 (40), 3.28 (50), 3.01 (60), 2.849 (60).

IMA No. 92-006

The nickel-analogue of hydromagnesite.

$\text{Ni}_5(\text{CO}_3)_4(\text{OH})_2 \cdot 4\text{--}5 \text{H}_2\text{O}$

Monoclinic: $P2_1/c$

a 10.06, b 8.75, c 8.32 Å, β 114.3°

Bluish-green; silky; transparent.

Biaxial (sign unknown), α' 1.630, γ' 1.640, $2V$
unknown.

6.30 (5), 5.75 (10), 4.36 (4), 4.14 (3), 2.871 (4b),
2.458 (2b), 2.120 (3).

IMA No. 92-008

$\text{NaH}(\text{CO}_3)\text{H}_3(\text{BO}_3) \cdot 2 \text{H}_2\text{O}$

Monoclinic: $C2$

a 16.119, b 6.928, c 6.730 Å, β 100.46°

Colourless; vitreous; transparent.

Biaxial (–), α 1.351 (calc.), β 1.459, γ 1.486,
 $2V(\text{meas.})$ 50°.

6.36 (25), 4.203 (6), 3.464 (100), 3.173 (59),
2.608 (5), 1.731 (19).

IMA No. 92-010

A triclinic polymorph of 92-011.

$\text{Ca}_9\text{B}_{26}\text{O}_{34}(\text{OH})_{24}\text{Cl}_4 \cdot 13 \text{H}_2\text{O}$

Triclinic: $P1$

a 12.759, b 13.060, c 9.733 Å, α 102.14°,
 β 102.03°, γ 85.68°

Colourless to very pale yellow; vitreous;
translucent to transparent.

Biaxial (+), α 1.537, β 1.548, γ 1.570,
 $2V(\text{meas.})$ 77°, $2V(\text{calc.})$ 71°.

9.21 (70), 7.69 (100), 5.74 (60), 4.63 (40),
3.845 (35), 2.199 (30b).

IMA No. 92-011

A monoclinic polymorph of 92-010.

$\text{Ca}_9\text{B}_{26}\text{O}_{34}(\text{OH})_{24}\text{Cl}_4 \cdot 13 \text{H}_2\text{O}$

Monoclinic: $P2_1$

a 19.88, b 9.715, c 17.551 Å, β 114.85°

Colourless to very pale yellow; vitreous;
translucent to transparent.

Biaxial (+), α 1.542, β 1.545, γ 1.565,
 $2V(\text{meas.})$ 47°, $2V(\text{calc.})$ 43°.

9.03 (60), 8.56 (100), 6.62 (70), 6.14 (30b),
5.12 (30), 4.09 (30), 3.768 (30), 3.493 (30).

IMA No. 92-012

$\text{Ca}_2(\text{CaMn})(\text{SiO}_3\text{OH})_2(\text{OH})_2$

Orthorhombic: $Pbca$

a 9.398, b 9.139, c 10.535 Å

Colourless; vitreous; transparent.

Biaxial (+), α 1.634, β 1.640, γ 1.656,
 $2V(\text{meas.})$ 65°, $2V(\text{calc.})$ 63°.

4.18 (45), 3.231 (100), 2.846 (42), 2.789 (35),
2.391 (42), 2.042 (28).

IMA No. 92-013

The phosphate analogue of preisingerite and
schumacherite.

$\text{Bi}_3\text{O}(\text{OH})(\text{PO}_4)_2$

Triclinic: $P\bar{1}$

a 9.798, b 7.250, c 6.866 Å, α 88.28°,
 β 115.27°, γ 110.70°

White to pale pink, sometimes brown;
vitreous; transparent to translucent.

Mean index of refraction estimated from
reflectance data: 2.01 at 589 nm.

4.437 (46), 3.247 (87), 3.188 (100), 3.135 (95),
3.026 (75), 2.953 (47), 2.165 (41).

IMA No. 92-014

$\text{Na}_x\text{Ca}_y\text{Cu}_z(\text{Mg},\text{Fe}^{3+},\text{Al})_3(\text{AsO}_4)_3$, $x \sim 0.76$,
 $y \sim 0.42$, $z \sim 0.39$

Monoclinic: $C2/c$

a 11.882, b 12.760, c 6.647 Å, β 112.81°

Light blue; vitreous; translucent.

Biaxial (+), α 1.714, β 1.744, γ 1.783,
 $2V(\text{meas.})$ 60°, $2V(\text{calc.})$ 84°.

4.35 (40), 4.06 (50), 3.56 (40), 3.53 (40),
3.495 (60), 3.066 (40), 2.744 (140),
2.605 (40).

IMA No. 92-015

The ferric analogue of millosevichite.

$(\text{Fe},\text{Al})_2(\text{SO}_4)_3$

Hexagonal: $R\bar{3}$

a 8.14, c 21.99 Å

White to light brown; dull; transparent.

Uniaxial (sign unknown), n is between 1.555
and 1.625.

5.99 (28), 4.35 (23), 3.56 (100), 2.97 (20),
2.72 (20), 2.64 (11).

IMA No. 92-016

The phosphate analogue of arsenoclasite.

$\text{Mn}_5(\text{PO}_4)_2(\text{OH})_4$

Orthorhombic: $P2_12_12_1$

a 9.097, b 5.693, c 18.00 Å

Pale yellow, yellow, pale burnt orange;
adamantine; transparent.

- Biaxial (sign unknown), α' 1.74, γ' 1.76, 2V unknown.
2.900 (100), 2.853 (70), 2.802 (50), 2.702 (80), 2.022 (15), 1.608 (15).
- IMA No. 92-017
A member of the garnet group.
 $\text{Ca}_3(\text{Ti,Fe}^{2+}\text{Fe}^{3+})_2(\text{Si,Fe}^{3+})_3\text{O}_{12}$
Cubic: Ia3d
a 12.162 Å
Black; adamantine; opaque.
Isotropic, ω 1.955.
3.039 (72), 2.720 (100), 2.483 (51), 2.385 (21), 1.973 (24), 1.687 (26), 1.626 (56).
- IMA No. 92-018
 $\text{Ca}_2\text{Y}(\text{AsO}_4)(\text{WO}_4)_2$
Tetragonal: I4₁/a
a 5.135, c 33.882 Å
Creamy yellow; vitreous to adamantine; translucent.
Uniaxial (+), ω 1.874, ε 1.918.
4.674 (18), 3.059 (100), 2.571 (19), 1.901 (32), 1.818 (16), 1.674 (17), 1.562 (32).
- IMA No. 92-019
 $\text{C}_{14}\text{H}_{10}$
Monoclinic: P2₁
a 8.392, b 6.181, c 9.558 Å, β 98.48°
Colourless to greyish-white; vitreous to waxy; transparent.
Biaxial (+), $n_{\text{min.}}$ ~1.75, $n_{\text{max.}}$ ~1.95, 2V(meas.) ~90°.
9.434 (100), 4.941 (11), 4.724 (11), 4.546 (5), 4.028 (13), 3.371 (10).
- IMA No. 92-020
A member of the amphibole group.
 $(\text{Na,K})(\text{Ca,Na})_2(\text{Mg,Fe}^{3+},\text{Fe}^{2+})_5\text{Si}_8\text{O}_{22}(\text{F,OH,O})_2$
Monoclinic: C2/m
a 9.762, b 17.888, c 5.122 Å, β 102.25°
Blue green and green; vitreous; transparent.
Biaxial (-), α 1.618, β 1.624, γ 1.627, 2V(meas.) 71°, 2V(calc.) 70°.
9.9 (70), 3.69 (60), 3.34 (100), 3.18 (60), 3.13 (90), 2.82 (70), 1.98 (90), 1.439 (60).
- IMA No. 92-024
 CuBi_2O_4
Tetragonal: P4/ncc
a 8.511, c 5.823 Å
Black; metallic; opaque.
In reflected light: grey, weak anisotropism, weak but distinct bireflectance, pleochroic grey with a faint bluish tint and brownish grey. $R_{\text{max.}}$ & $R_{\text{min.}}$: (21.1, 19.0%) 482 nm, (20.2, 18.0%) 545 nm, (19.7, 17.6%) 589 nm, (19.5, 17.3%) 659 nm.
- 4.26 (17), 3.191 (100), 2.913 (16), 2.695 (18), 1.947 (18).
- IMA No. 92-025
 $\text{Cu}_3\text{TeO}_6 \cdot \text{H}_2\text{O}$
Cubic: P-lattice, space group unknown
a 9.555 Å
Emerald green; adamantine; transparent to translucent.
Isotropic, ω 2.01 calculated from reflectance values at 589 nm.
4.26 (40), 2.763 (100), 2.384 (70), 1.873 (40), 1.689 (80), 1.440 (60).
- IMA No. 92-026
The -2H polytype of 92-027.
 $\text{Mn}_4\text{Al}_2(\text{OH})_{12}\text{CO}_3 \cdot 3\text{H}_2\text{O}$
Hexagonal: P6₃22
a 10.985, c 15.10 Å
Orange-brown, pale brown, pale blue, colourless; vitreous; transparent.
Uniaxial (-), ω 1.587, ε 1.547.
7.53 (100), 3.768 (60), 2.578 (50), 2.221 (40), 1.856 (40), 1.552 (40).
- IMA No. 92-027
The -3T polytype of 92-026.
 $\text{Mn}_4\text{Al}_2(\text{OH})_{12}\text{CO}_3 \cdot 3\text{H}_2\text{O}$
Hexagonal (trigonal): P3₁12 or P3₂12
a 10.985, c 22.63 Å
Orange-brown, pale brown; vitreous; transparent.
Uniaxial (-), ω 1.587, ε could not be measured.
7.55 (100), 3.770 (90), 2.670 (70), 2.346 (70), 1.973 (60), 1.586 (30), 1.662 (30).
- IMA No. 92-028
The -2H polytype of 92-029.
 $\text{Mg}_4\text{Al}_2(\text{OH})_{12}\text{CO}_3 \cdot 3\text{H}_2\text{O}$
Hexagonal: P6₃22
a 10.571, c 15.139 Å
Orange-brown, pale brown; vitreous; transparent.
Uniaxial (+), ω 1.533, ε 1.533.
7.63 (100), 3.785 (100), 2.603 (15), 2.496 (15), 2.341 (15), 2.166 (15), 1.991 (15), 1.825 (20), 1.495 (15).
- IMA No. 92-029
The -3T polytype of 92-028.
 $\text{Mg}_4\text{Al}_2(\text{OH})_{12}\text{CO}_3 \cdot 3\text{H}_2\text{O}$
Hexagonal (trigonal): P3₁12 or P3₂12
a 10.558, c 22.71 Å
Yellow to pale yellow; vitreous; transparent.
Uniaxial (+ or -), ω 1.533, ε 1.533.
7.57 (100), 3.778 (90), 2.570 (40), 2.281 (40), 1.932 (40), 1.524 (20), 1.493 (20).

IMA No. 92-030

$\text{Fe}_4\text{Al}_2(\text{OH})_{12}\text{CO}_3 \cdot 3\text{H}_2\text{O}$
Hexagonal (trigonal): P3_112 or P3_212
a 10.805, c 22.48 Å
Green-brown with black coating; vitreous;
transparent.
Uniaxial (-), ω 1.599, ϵ 1.570.
7.49 (100), 3.746 (50), 2.625 (40), 2.314 (50),
1.948 (40), 1.558 (15), 1.526 (20).

IMA No. 92-031

$\text{Na}_5\text{YZrSi}_6\text{O}_{18} \cdot 6\text{H}_2\text{O}$
Hexagonal (trigonal): R32
a 10.825, c 15.809 Å
Light green to yellow green; vitreous;
transparent to translucent.
Uniaxial (-), ω 1.585, ϵ 1.578.
6.03 (32), 5.40 (63), 3.236 (84), 3.127 (88),
3.030 (100), 1.805 (21).

IMA No. 92-032

A member of the amphibole group.
 $(\text{K},\text{Na})(\text{Na},\text{Li})_2(\text{Mg},\text{Mn}^{3+},\text{Fe}^{3+},\text{Li})_5\text{Si}_8\text{O}_{22}(\text{OH})_2$
Monoclinic: $\text{P2}_1/\text{m}$
a 9.94, b 17.80, c 5.302 Å, β 105.5°
Dark red to brownish lilac; vitreous;
transparent.
Biaxial (-), α 1.654, β 1.675 (calculated),
 γ 1.696, 2V(meas.) 88–92°.
8.890 (M), 8.427 (M), 5.077 (M), 4.442 (M),
3.357 (M), 3.257 (S), 3.132 (S), 2.812 (S),
2.553 (S) plus seven other lines of intensity
(M).

IMA No. 92-033

$\text{SrMn}_2^{3+}[\text{Si}_2\text{O}_7](\text{OH})_2 \cdot \text{H}_2\text{O}$
Orthorhombic: Cmcm
a 6.245, b 9.031, c 13.404 Å
Orange-brown; vitreous; translucent.
Biaxial (+), n's > 1.82, 2V(meas.) 63°.
4.804 (86), 3.373 (66), 2.833 (100), 2.807 (82),
2.695 (98), 2.401 (68).

IMA No. 92-034

A member of the tourmaline group.
 $\square(\text{Fe}_2^+\text{Al})\text{Al}_6\text{Si}_6\text{O}_{18}(\text{BO}_3)_3(\text{OH})_4$
Hexagonal (trigonal): R3m
a 15.967, c 7.126 Å
Bluish black; vitreous; transparent.
Uniaxial (-), ω 1.664, ϵ 1.642.
6.338 (84), 4.212 (48), 3.989 (38), 3.452 (91),
2.944 (71), 2.573 (100).

IMA No. 92-035

The magnesium-analogue of staurolite.
 $(\text{Mg},\text{Li},\text{Fe},\square)_4\text{Al}_{18}\text{Si}_8\text{O}_{44}(\text{OH})_4$
Monoclinic: $\text{C2}/\text{m}$

a 7.872, b 16.55, c 5.634 Å, β 90.00°
Colourless in thin section; vitreous to
resinous; transparent.
Biaxial (sign unknown), mean n 1.709,
2V unknown.
4.139 (24), 2.678 (38), 2.390 (50), 2.370 (33),
2.356 (24), 1.968 (100).

IMA No. 92-036

The zinc-analogue of staurolite.
 $(\text{Zn},\text{Li},\text{Fe},\text{Mg},\square)_4\text{Al}_{18}\text{Si}_8\text{O}_{44}(\text{OH})_4$
Monoclinic: $\text{C2}/\text{m}$
a 7.853, b 16.54, c 5.639 Å, β 90.00°
Colourless in thin section; vitreous to
resinous; transparent.
Biaxial (sign unknown), $\alpha \sim 1.722$,
 β unknown, γ 1.734, 2V unknown.
3.001 (61), 2.678 (70), 2.390 (87), 2.363 (46),
2.349 (45), 1.968 (61), 1.964 (48),
1.391 (100).

IMA No. 92-037

The tetragonal, lead-analogue of lavendulan.
 $\text{NaPbCu}_5(\text{AsO}_4)_4\text{Cl} \cdot 5\text{H}_2\text{O}$
Tetragonal: P4_122 or P4_322
a 10.066, c 39.39 Å
Intense blue; vitreous; translucent.
Uniaxial (-), ω 1.770, ϵ 1.710.
9.83 (100), 4.925 (60), 4.482 (50), 3.132 (90),
2.772 (40), 2.515 (50), 1.778 (40).

IMA No. 92-038

$\text{Cu}_2\text{O}(\text{Fe},\text{Cu},\text{Zn})_6\text{Mo}_2\text{Ge}_6\text{S}_{32}$
Cubic: space group unknown
a 10.64 Å
Megascopic colour unknown; metallic;
opaque.
In reflected light: pale yellow to greyish
yellow, no anisotropism, no bireflectance,
nonpleochroic. R: (23.7%) 470 nm, (25.5%)
546 nm, (25.7%) 589 nm, (25.6%) 650 nm.
3.07 (10), 2.66 (2), 1.884 (8), 1.603 (4),
1.536 ($1/2$), 1.331 (1), 1.220 (2), 1.190 (1).

IMA No. 92-039

$\text{Cu}_{20}(\text{Fe},\text{Zn},\text{Cu})_6\text{W}_2\text{Ge}_6\text{S}_{32}$
Cubic: space group unknown
a 10.675 Å
Megascopic colour unknown; metallic;
opaque.
In reflected light: pale yellowish pink, no an-
isotropism, no bireflectance, nonpleochroic.
R: (23.2%) 470 nm, (23.7%) 546 nm,
(24.0%) 589 nm, (23.8%) 650 nm.
4.36 (1), 3.38 (1), 3.08 (10), 2.67 (2), 1.887 (7),
1.612 (5), 1.543 (1), 1.333 (1), 1.225 ($1/2$),
1.192 ($1/2$).

- IMA No. 92-040
 $\text{Na}_4\text{Zn}_2\text{Si}_7\text{O}_{18} \cdot 5 \text{H}_2\text{O}$
 Orthorhombic: F2dd
 a 10.211, b 39.88, c 10.304 Å
 Colourless to light mauve; vitreous;
 transparent.
 Biaxial (+), α 1.520, β 1.521, γ 1.524,
 2V(meas.) 61°, 2V(calc.) 60°.
 6.346 (10), 4.959 (3), 3.240 (6), 3.167 (4),
 3.140 (4), 2.821 (3).
- IMA No. 92-041
 The thallium-analogue of jarosite.
 $(\text{Tl},\text{K})\text{Fe}_3(\text{SO}_4)_2(\text{OH})_6$
 Hexagonal (trigonal): R3m
 a 7.3301, c 17.6631 Å
 Gold yellow; adamantine; transparent.
 Uniaxial (-), ω 1.822, ε 1.768.
 5.974 (87), 3.666 (34), 3.112 (100), 2.9877 (22),
 2.5773 (21), 1.9912 (29), 1.8329 (23).
- IMA No. 92-043
 $\text{Ca}(\text{UO}_2)_4(\text{SO}_4)_2(\text{OH})_6 \cdot 6 \text{H}_2\text{O}$
 Orthorhombic: P-lattice, space group
 unknown
 a 8.73, b 17.09, c 15.72 Å
 Sulphur yellow; vitreous; translucent.
 Biaxial (-), α 1.617 (calculated), β 1.710,
 γ 1.758, 2V(meas.) 68°.
 7.90 (100), 4.17 (30), 3.98 (40), 3.49 (80),
 3.38 (70), 2.844 (30b).
- IMA No. 92-045
 The phosphate-analogue of segnitite.
 $\text{PbFe}_3^{3+}(\text{PO}_4)_2(\text{OH},\text{H}_2\text{O})_6$
 Hexagonal (trigonal): R3m
 a 7.325, c 16.900 Å
 Cream to brownish yellow to yellowish green;
 adamantine; translucent.
 Uniaxial (-), ω 1.955, ε 1.935.
 5.96 (90), 3.67 (60), 3.07 (100), 2.538 (50),
 2.257 (50), 1.979 (50).
- IMA No. 92-046
 $\text{AlF}_3 \cdot 3 \text{H}_2\text{O}$
 Tetragonal: P4/n
 a 7.715, c 3.648 Å
 Colourless; vitreous; transparent.
 Uniaxial (-), ω 1.427, ε 1.403.
 5.47 (100), 2.439 (72), 2.027 (70), 1.775 (78),
 1.725 (85), 1.306 (70).
- IMA No. 92-048
 $\text{Na}_4\text{REE}_2(\text{CO}_3)_5$ with Ce the dominant REE
 Monoclinic: P2₁
 a 20.84, b 6.374, c 10.578 Å, β 120.45
- Grey with slight pinkish tint; vitreous;
 translucent.
 Biaxial (+ or -), α 1.623, β 1.636, γ 1.649,
 2V(meas.) 90°, 2V(calc.) 89°.
 9.13 (3), 5.22 (5), 4.13 (3), 3.70 (4), 2.607 (10),
 2.148 (3), 1.921 (3).
- IMA No. 92-050
 The magnesium-analogue of dumortierite.
 $(\text{Mg},\text{Ti},\square)(\text{Al},\text{Mg})_2\text{Al}_4\text{Si}_3\text{O}_{18-x}(\text{OH})_xB$ $x \approx 3$
 Orthorhombic: Pmcn
 a 12.02, b 20.22, c 4.732 Å
 Pink to red; vitreous; transparent.
 Biaxial (-), α 1.678, β 1.700, γ 1.701,
 2V(meas.) 38°, 2V(calc.) 24°.
 6.01 (59), 5.88 (100), 3.489 (60), 3.255 (82),
 3.074 (53), 2.927 (74), 2.131 (50), 2.090 (48).
- NOTE:**
- The following three minerals from previous years
 also have been approved.
- IMA No. 90-006
 $\text{Fe}_{16}\text{O}_{16}(\text{OH})_y(\text{SO}_4)_z$
 where $16 - y = 2z$ and $2.0 \leq z \leq 3.5$
 Tetragonal: probably P4/m
 a 10.66, c 6.04 Å
 Brownish yellow; dull; translucent.
 Optical properties unknown.
 4.86 (37), 3.38 (46), 2.55 (100), 2.28 (23),
 1.66 (21), 1.51 (24), 1.46 (18).
- IMA No. 90-046
 The uranium-analogue of polycrase-(Y).
 $(\text{U},\text{Y})(\text{Ti},\text{Nb},\text{Ta})_2\text{O}_8$
 Orthorhombic: Pbcn
 a 14.48, b 5.559, c 5.223 Å
 Brown-red; adamantine; opaque.
 In reflected light: pale grey with bluish tones;
 no anisotropism, bireflectance, or
 pleochroism. R: (23.6%) 470 nm, (21.5%)
 546 nm, (22.3%) 589 nm, (25.1%) 650 nm.
 3.73 (W), 3.21 (W), 2.99 (S), 2.78 (W),
 1.90 (MS), 1.86 (W), 1.77 (MW), 1.48 (M).
- IMA No. 91-036
 $\text{Fe}_2(\text{OH})_3\text{Cl}$
 Orthorhombic: Pnam
 a 6.31, b 9.20, c 7.10 Å
 Megascopic colour unknown; lustre probably
 dull; transparent.
 Index of refraction: 1.6 to 1.7.
 Electron diffraction pattern: 5.68, 5.07, 2.93,
 2.37, 2.14, 1.65.