

Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =
Bulletin suisse de minéralogie et pétrographie

Band: 77 (1997)

Heft: 2

Vereinsnachrichten: New minerals recently approved by the Commission on New
Minerals and Mineral Names : International Mineralogical
Association

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 17.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

**New minerals recently approved
by the
Commission on New Minerals and Mineral Names
International Mineralogical Association**

The information given here is provided by the Commission on New Minerals and Mineral Names, I. M. A. for comparative purposes and as a service to mineralogists working on new species.

Each mineral is described in the following format:

IMA No. (any relationship to other minerals)
Chemical Formula
Crystal system, space group
unit cell parameters
Colour; lustre; diaphaneity
Optical properties
Strongest lines in the X-ray powder diffraction pattern


The names of these approved species are considered confidential information until the authors have published their descriptions or released information themselves.

No other information will be released by the commission.

J. A. Mandarino, Chairman Emeritus, and J. D. Grice, Chairman
Commission on New Minerals and Mineral Names, International Mineralogical Association

1996 Proposals

IMA No. 96-001



Triclinic: P1 or P $\bar{1}$

a 8.984, b 10.079, c 8.975 Å, α 102.68°,
 β 92.45°, γ 70.45°


Emerald green; vitreous to adamantine;
transparent to translucent

Biaxial, indices of refraction calculated from
reflectance measurements are 1.71–1.73

9.28 (70), 4.65 (70), 3.097 (100), 3.018 (60),
2.658 (50), 2.468 (50), 1.740 (50)

IMA No. 96-002

The calcium-dominant analogue of guillemite



Orthorhombic: Pmn2₁ or Pmnm

a 7.010, b 17.135, c 17.606 Å

Lemon-yellow; pearly; translucent

Biaxial (–), α 1.54 calc., β 1.73, γ 1.75,
2V(meas.) 33°

8.79 (80), 8.56 (40), 3.51 (100), 3.24 (40),
3.093 (50), 3.032 (100), 1.924 (40)

IMA No. 96-003

The Fe³⁺-dominant analogue of winstanleyite
(Fe³⁺, Te⁶⁺, Ti⁴⁺, Mg²⁺)Te₃O₈

Cubic: Ia $\bar{3}$

a 11.011 Å

Orange; adamantine; translucent

Isotropic, n(calc) = 2.17


4.486 (29), 3.175 (100), 2.943 (23), 2.749 (37),
2.592 (22), 1.944 (44), 1.658 (45)

- IMA No. 96-004
 $\text{Al}_2(\text{OH})_5\text{Cl} \cdot 2\text{H}_2\text{O}$
 Cubic: $\text{Im}\bar{3}\text{m}$
 a 19.878 Å
 Yellow-orange to yellow-brown; vitreous;
 transparent
 Isotropic, n 1.53–1.55
 8.11 (70), 7.03 (50), 4.47 (60), 3.23 (70),
 2.706 (100), 2.446 (80), 1.957 (70)
- IMA No. 96-005
 $\text{Mg}_2(\text{CO}_3)\text{Cl}(\text{OH}) \cdot 3\text{H}_2\text{O}$
 Hexagonal (trigonal): $\text{R}\bar{3}\text{c}$ or $\text{R}\bar{3}\text{c}$
 a 23.163, c 7.221 Å
 White; lustre and diaphaneity unknown
 Uniaxial, ω 1.510, ε 1.510
 11.66 (100), 3.396 (17), 3.356 (17), 3.264 (21),
 3.218 (21), 3.000 (41), 2.657 (22)
- IMA No. 96-006
 $\text{NaZn}_4(\text{SO}_4)(\text{OH})_6\text{Cl} \cdot 6\text{H}_2\text{O}$
 Hexagonal (trigonal): $\text{P}\bar{3}$
 a 8.359, c 13.059 Å
 Colourless to white; pearly; translucent
 Uniaxial (–), ω 1.5607, ε 1.5382
 14.244 (100), 6.501 (23), 4.339 (15), 3.258 (14),
 2.967 (10)
- IMA No. 96-007
 $(\text{K},\text{Na})_5\text{Fe}_7^{3+}[\text{Si}_{20}\text{O}_{50}(\text{OH})_6] \cdot 12\text{H}_2\text{O}$
 Triclinic: $\text{P}\bar{1}$
 a 14.86, b 20.54, c 5.29 Å, α 95.6°, β 92.3°,
 γ 94.4°
 Pink-brownish; silky to earthy; translucent
 Biaxial (+), α 1.523, β 1.525, γ 1.550,
 2V(meas.) 30°, 2V(calc.) 32°
 12.36 (100), 11.60 (40), 10.21 (14), 3.411 (37),
 3.281 (15), 2.896 (12)
- IMA No. 96-008
 The Fe^{3+} -dominant analogue of beryl
 $(\text{Fe},\text{Mg},\text{Al})_2(\text{Na},\square)[\text{Be}_3\text{Si}_6\text{O}_{18}] \cdot \text{H}_2\text{O}$
 Hexagonal: $\text{P}6/\text{mcc}$
 a 9.387, c 9.202 Å
 Light-blue; vitreous; transparent
 Uniaxial (–), ω 1.625, ε 1.619
 8.12 (S), 4.00 (M), 3.278 (VS), 2.903 (S),
 2.553 (MW), 1.752 (MW)
- IMA No. 96-009
 $\text{Ca}_3[\text{B}_5\text{O}_6(\text{OH})_7]\text{Cl}_2 \cdot 8\text{H}_2\text{O}$
 Monoclinic: Pn
 a 17.42, b 8.077, c 17.33 Å, β 121.48°
 Colourless to white; vitreous; transparent
 to translucent
 Biaxial (–), α 1.506, β 1.527, γ 1.532,
 2V(meas.) 56°, 2V(calc.) 51°
- 8.10 (10), 4.04 (4), 3.56 (2), 2.834 (2),
 2.535 (2), 2.276 (2)
- IMA No. 96-010
 The Fe^{3+} -dominant analogue of tomichite
 $(\text{Fe}^{3+},\text{Ti})_4\text{Ti}_3\text{AsO}_{13}(\text{OH})$
 Monoclinic: $\text{A}2/\text{m}$
 a 7.184, b 14.289, c 5.006 Å, β 105.17°
 Black; metallic; opaque
 In reflected light: greyish-white, no bire-
 flectance, nonpleochroic. R_1 & R_2 :
 (20.1, 20.8%) 460 nm, (18.7, 19.3%) 540 nm,
 (18.2, 18.9%) 580 nm, (17.5, 18.1%) 660 nm
 3.117 (30), 2.846 (80), 2.681 (100),
 2.029 (30), 1.5825 (50)
- IMA No. 96-012
 $\text{Ca}(\text{H}_2\text{O})_3(\text{C}_2\text{O}_4)$ or $\text{CaC}_2\text{O}_4 \cdot 3\text{H}_2\text{O}$
 Triclinic: $\text{P}\bar{1}$
 a 6.097, b 7.145, c 8.434 Å, α 76.54°,
 β 70.30°, γ 70.75°
 Colourless; vitreous; transparent
 Biaxial (–), α' 1.483, β 1.516(calc.), γ' 1.533,
 2V(meas.) 70°, 2V(calc.) 70°
 7.92 (M), 5.52 (VS), 5.26 (M), 4.99 (M),
 3.642 (M), 2.834 (S), 2.758 (M), 2.732 (M)
- IMA No. 96-013
 $\text{Fe}^{2+}(\text{UO}_2)_2(\text{SO}_4)_2(\text{OH})_2 \cdot 3\text{H}_2\text{O}$
 Orthorhombic: Pnmm or $\text{Pnnm}2$
 a 15.908, b 16.274, c 6.903 Å
 Pale yellow to white; vitreous; transparent
 Biaxial (–), α 1.470, β 1.492, γ 1.504(calc.),
 2V(meas.) 73°
 7.95 (81), 5.91 (100), 3.941 (71), 3.451 (67),
 3.166 (50), 2.894 (41), 2.596 (70)
- IMA No. 96-014
 $\text{Pb}_{14}\text{Sb}_{30}\text{S}_{54}\text{O}_{55}$
 Monoclinic: $\text{C}2/\text{m}$
 a 52.00, b 8.148, c 24.311 Å, β 104.09°
 Bluish-black; metallic; opaque
 In reflected light: black with blue-red reflec-
 tions, low anisotropism, low bireflectance,
 nonpleochroic. R_1 & R_2 : (40.03, 42.90%)
 470 nm, (36.46, 40.92%) 546 nm, (35.65,
 40.25%) 589 nm, (32.40, 36.00%) 650 nm
 4.04 (m), 3.47 (s), 3.44 (m), 3.04 (m), 2.96 (s),
 2.296 (m)
- IMA No. 96-015
 $\text{Cu}_5\text{O}_2(\text{SeO}_3)_2\text{Cl}_2$
 Monoclinic: $\text{P}2_1/\text{c}$
 a 6.045, b 13.778, c 5.579 Å, β 95.76°
 Chestnut to dark brown; very strong vitreous
 to adamantine; translucent

- Biaxial (–), α 2.06, β 2.11, γ 2.15, 2V(meas.) large, 2V(calc.) 82°
6.88 (68), 5.511 (50), 2.990 (100), 2.963 (94), 2.566 (67), 2.296 (95)
- IMA No. 96-017
A triclinic polymorph of clinoclase
 $\text{Cu}_3(\text{AsO}_4)(\text{OH})_3$
Triclinic: P1
a 5.445, b 5.873, c 5.104 Å, α 114.95°, β 93.05°, γ 91.92°
Green-blue; vitreous; transparent
Biaxial (–), α 1.760, β 1.80, γ 1.83, 2V(meas.) 77° , 2V(calc.) 80°
4.613 (100), 4.580 (50), 3.390 (60), 2.713 (40), 2.543 (40), 2.445 (30)
- IMA No. 96-019
 $\text{NaCa}_3(\text{CO}_3)_2\text{F}_3 \cdot \text{H}_2\text{O}$
Hexagonal (trigonal): P3₂
a 6.718, c 15.050 Å
Colourless to white; vitreous; transparent to translucent
Uniaxial (+), ω 1.538, ε 1.563
5.809 (30), 5.010 (30), 3.358 (30), 2.791 (50), 2.508 (40), 2.010 (100), 1.939 (40)
- IMA No. 96-020
 $\text{Pb}_{12}\text{O}_6\text{Mn}(\text{Mn},\text{Mg})_4(\text{Mg},\text{Mn})_2(\text{SO}_4)(\text{CO}_3)_4\text{Cl}_4(\text{OH})_{12}$
Tetragonal: P4₂/nnm
a 12.627, c 12.595 Å
Apple green to emerald green; vitreous to adamantine; transparent
Anomalously biaxial (+), α , β , and $\gamma > 1.92$
8.95 (20), 7.30 (20), 3.99 (30), 2.975 (100), 2.752 (30), 2.473 (20), 1.716 (20)
- IMA No. 96-022
A polymorph of fluorapatite
 $(\text{Ca},\text{R})_5(\text{PO}_4)_3\text{F}$ R = Sr, Na, REE
Hexagonal: P6₃
a 9.485, c 7.000 Å
Pale yellow; vitreous; transparent
Uniaxial (–), ω 1.649, ε 1.637
3.498 (45), 3.104 (22), 2.838 (100), 2.814 (48), 2.740 (53), 1.963 (21), 1.865 (31)
- IMA No. 96-023
A manganese- and fluorine-rich member of the eudialyte group
 $(\text{Na},\text{RE})_{15}(\text{Ca},\text{RE})_6\text{Mn}_3\text{Zr}_3\text{NbSi}_{25}\text{O}_{76}\text{F}_2$
Hexagonal (trigonal): R3m
a 14.1686, c 30.0847 Å
Yellow-brown; vitreous; transparent
Uniaxial (–), ω 1.628, ε 1.623
- 11.362 (43), 7.084 (41), 5.681 (30), 4.296 (34), 3.382 (37), 2.962 (91), 2.840 (100)
- IMA No. 96-024
The scandium-dominant analogue of xenotime-(Y)
 ScPO_4
Tetragonal: I4₁/amd
a 6.589, c 5.806 Å
Pale-pink; vitreous; transparent
Uniaxial (+), ω 1.790, ε 1.86
3.293 (100), 2.464 (8), 2.178 (4), 2.055 (4), 1.693 (6), 1.647 (6)
- IMA No. 96-025
A member of the zeolite group
 $\text{Na}_3\text{Ca}_4\text{Al}_{11}\text{Si}_{85}\text{O}_{192} \cdot 60\text{H}_2\text{O}$
Orthorhombic: Pnma
a 20.223, b 20.052, c 13.491 Å
Colourless to milky-white; silky to vitreous; opaque to transparent
Biaxial (–), α 1.485, β 1.487, γ 1.488, 2V(calc.) 70°
11.20 (84), 9.98 (35), 3.85 (100), 3.75 (98), 3.67 (27), 3.00 (32)
- IMA No. 96-026
An orthorhombic polymorph of corderoite
 $\gamma\text{-Hg}_3\text{S}_2\text{Cl}_2$
Orthorhombic: Ammm, A222 or A2mm (Am2m, Amm2)
a 9.332, b 16.82, c 9.108 Å
Canary yellow; glassy; transparent
Biaxial (+), mean index of refraction 2.25, 2V(meas.) $> 70^\circ$
In reflected light: white, anisotropism and bireflectance not observed, R(est.) $\approx 15\%$
3.65 (90), 3.11 (51), 2.83 (36), 2.60 (49), 2.58 (100), 2.33 (41), 2.11 (31)
- IMA No. 96-027
 $\text{NaCu}_5\text{O}_2(\text{SeO}_3)_2\text{Cl}_3$
Orthorhombic: Pbnm
a 10.482, b 17.732, c 6.432 Å
Emerald-green; vitreous; transparent
Biaxial (–), α 1.845, β 1.968, γ 1.975, 2V(meas.) 20° , 2V(calc.) 31°
9.01 (10), 8.84 (60), 5.24 (100), 3.251 (40), 2.955 (27), 2.626 (25), 2.513 (12)
- IMA No. 96-028
 $\text{NaFe}_4^{2+}(\text{PO}_4)_3$
Hexagonal (trigonal): R3
a 14.97, c 41.66 Å
Very pale amber; waxy; transparent
Uniaxial (+), ω 1.72, ε 1.75

4.13 (80), 3.47 (50), 3.21 (50), 3.01 (90),
2.93 (50), 2.85 (50), 2.71 (100), 2.57 (50)

IMA No. 96-029



Monoclinic: C2/m, Cm or C2


a 14.767, b 5.574, c 15.079 Å, β 91.959°

White; vitreous; transparent

Biaxial (+), α 1.629, β 1.640, γ 1.654,
2V(meas.) 82°, 2V(calc.) 84°

5.41 (70), 5.19 (100), 4.95 (60), 4.31 (70),
3.378 (60), 2.162 (40)

IMA No. 96-030




Orthorhombic: Pnma or Pn2₁a

a 5.866, b 3.893, c 7.302 Å

Colour not observed, metallic, opaque

In reflected light: brownish with a pale green
tinge, anisotropism moderate-distinct from
dark brown to pale greyish green, bire-
flectance weak, pleochroism brownish to
greenish. R_{min} & R_{max} : (45.5, 46.3%)
470 nm, (47.6, 48.4%) 546 nm, (48.2,
49.5%) 589 nm, (49.8, 51.2%) 650 nm
2.426 (7), 2.348 (4), 2.237 (10), 2.067 (8),
1.935 (6), 1.860 (5)

IMA No. 96-032



Orthorhombic: Pmcn


a 6.964, b 9.173, c 6.302 Å

Colourless to pale yellow; vitreous;
transparent and translucent

Biaxial (–), α 1.457, β 1.543, γ 1.622,
2V(meas.) 82°, 2V(calc.) 83°

5.19 (90), 3.477 (100), 2.800 (50), 2.087 (50),
2.057 (50), 1.966 (50), 1.849 (50), 1.763 (50)

IMA No. 96-033



Hexagonal: P6₃/mmc

a 5.068, c 22.87 Å


Colourless to slightly beige; vitreous to some-
what pearly; transparent to translucent

Uniaxial (–), ω 1.728, ε 1.542

4.31 (100), 3.169 (70), 2.877 (60), 2.534 (70),
2.192 (90B), 1.978 (70)

IMA No. 96-034

The magnesium- and phosphate-dominant
analogue of allactite



Monoclinic: P2₁/n

a 5.250, b 11.647, c 9.655 Å, β 95.93°


Colourless; pearly; transparent

Biaxial (–), α 1.5945, β 1.6069, γ 1.6088,
2V(meas.) 46°, 2V(calc.) 43°

4.436 (75b), 3.521 (80), 3.145 (70), 3.087 (70),
2.905 (100), 2.794 (75), 2.199 (80)

IMA No. 96-035

The phosphate-dominant analogue
of gartrellite



Triclinic: P1 or P1̄

a 5.320, b 5.528, c 7.434 Å, α 67.61°,
 β 69.68°, γ 70.65°


Green; vitreous to adamantine; transparent
to translucent

Biaxial (+), α 1.90, β 1.93 (calc.), γ 2.00,
2V(meas.) 70°

4.720 (67), 4.502 (61), 4.360 (100), 3.250 (70),
3.138 (57), 2.885 (89), 2.868 (69)

IMA No. 96-036

The calcium-dominant analogue of steacyite



Tetragonal: P4/mcc

a 7.592, b 7.592, c 14.824 Å

Apple-green to dark-green and brown;
vitreous; transparent

Uniaxial (–), ω 1.611, ε 1.606

5.36 (40), 5.31 (70), 3.40 (100), 3.33 (65),
2.654 (59), 2.231 (50)

IMA No. 96-037



Cubic: I4̄3m

a 15.470 Å


Pale greenish blue; vitreous; transparent

Isotropic, n 1.566

10.8 (29), 7.73 (34), 3.164 (100), 2.827 (28),
2.738 (29), 2.582 (37), 2.445 (36)

IMA No. 96-038

The copper-dominant analogue of carrboy-
dite and glaucocerinite



Hexagonal (trigonal): R3̄m

a 3.070, c 31.9 Å


Blue to pale blue; vitreous; translucent

Uniaxial (+), n_{min} 1.549, n_{max} 1.565

10.5 (100), 5.26 (17), 3.50 (6), 2.60 (5),
2.46 (2), 2.23 (2), 1.524 (5b)

IMA No. 96-039

The chromium-dominant analogue
of haxonite



Cubic: Fm3m

a 10.65 Å

- Iron-grey; metallic; opaque
In reflected light: white. R: (46.5%) 470 nm, (43.7%) 546 nm, (43.2%) 589 nm, (44.4%) 660 nm
2.38 (3), 2.17 (5), 2.05 (10)
- IMA No. 96-040
The antimony-dominant analogue of calzirtite
 $\text{Ca}_2(\text{Zr}, \text{Ti})_5(\text{Sb}^{5+}, \text{Mn}^{3+})_2\text{O}_{16}$
Tetragonal: $I4_1/acd$
a 15.199, c 10.181 Å
Bright red; adamantine; translucent
Uniaxial (+), ω 2.12, ε 2.16
3.45 (40), 2.92 (100), 2.539 (60), 1.794 (90), 1.535 (80), 1.0353 (40)
- IMA No. 96-041
The titanium-dominant analogue of brannockite
 $\text{KLi}_3\text{Ti}_2\text{Si}_{12}\text{O}_{30}$
Hexagonal: $P6/mcc$
a 9.903, c 14.276 Å
White; vitreous; transparent
Uniaxial (-), ω 1.635, ε 1.630
7.15 (40), 4.29 (50), 4.07 (85), 3.57 (80), 3.16 (100), 2.895 (95), 2.742 (30)
- IMA No. 96-043
The antimony-dominant analogue of fleischerite (with AsO_4 replacing one SO_4)
 $\text{Pb}_3\text{Sb}^{5+}(\text{SO}_4)(\text{AsO}_4)(\text{OH})_6 \cdot 3\text{H}_2\text{O}$
Hexagonal: $P6_322$
a 8.939, c 11.102 Å
Colourless; adamantine; transparent
Uniaxial (+), ω 1.760, ε 1.801
6.35 (44), 3.655 (100), 3.481 (80), 3.175 (31), 2.675 (62), 2.235 (35)
- IMA No. 96-044
 $\text{Ag}_2\text{Pd}_3\text{Se}_4$
Monoclinic: $P2_1/m$ or $P2_1$
a 6.350, b 10.387, c 5.683 Å, β 114.90°
Colour unknown, only visible in polished section; metallic; opaque
In reflected light: buff to slightly grey-green buff; moderate anisotropism, rotation tints rose-brown, grey-green, pale bluish grey and dark steel-blue; bireflectance weak (air), moderate (oil); very weak pleochroism. $R_1, R_2; {}^{im}R_1, {}^{im}R_2$: (39.7, 47.2; 26.2, 34.4%) 470 nm, (43.1, 48.8; 29.3, 35.15%) 546 nm, (44.3, 49.4; 30.4, 35.5%) 589 nm, (44.4, 49.2; 31.0, 35.6%) 650 nm
2.868 (50b), 2.742 (100), 2.688 (80), 2.367 (50), 1.956 (100), 1.829 (30)
- IMA No. 96-045
 $\text{Pb}_{7.5}\text{B}_{0.5}(\text{OH})_{3.5}\text{O}_{4.5}\text{Cl}_4$ or $\text{Pb}_8\text{O}_4(\text{OH})_4\text{Cl}_4$
Monoclinic: $C2/c$
a 5.673, b 5.580, c 13.152 Å, β 90.47°
Pale yellow to reddish orange; vitreous, resinous; translucent
In reflected light: grey; internal reflections ubiquitous, amber to light yellow; anisotropism masked (if present) by the internal reflections; bireflectance weak, non-pleochroic. $R_1, R_2; {}^{im}R_1, {}^{im}R_2$: (15.2, 16.3; 4.07, 4.67%) 470 nm, (14.2, 15.3; 3.59, 4.17%) 546 nm, (13.9, 15.0; 3.44, 4.02%) 589 nm, (13.7, 14.7; 3.37, 3.91%) 650 nm
6.581 (37), 3.785 (48), 3.267 (35), 2.930 (100), 2.825 (43), 2.780 (36), 2.182 (37), 1.980 (33)
- IMA No. 96-047
The iron-dominant analogue of cuprorhodsite
 $(\text{Fe}, \text{Cu})(\text{Rh}, \text{Ir}, \text{Pt})_2\text{S}_4$
Cubic: $Fd3m$
a 9.89 Å
Black; metallic; opaque
In reflected light: white, isotropic. R: (41.4%) 470 nm, (41.8%) 546 nm, (41.8%) 589 nm, (41.7%) 650 nm
5.72 (7), 2.99 (10), 2.471 (8), 1.903 (7), 1.750 (9), 1.674 (3), 1.009 (3)
- IMA No. 96-048
 $\text{Cu}_9\text{O}_2(\text{SeO}_3)_4\text{Cl}_6$
Monoclinic: $I2$
a 14.110, b 6.27, c 12.997 Å, β = 113.0°
Tobacco-green; strong vitreous; transparent
Biaxial (-), α 1.87, β 1.92, γ 1.94,
2V(meas.) 66°, 2V(calc.) 63°
11.29 (63), 5.56 (83), 3.450 (100), 3.239 (39), 2.714 (33), 2.486 (61)
- IMA No. 96-049
 $\text{CaMgNa}_6(\text{IO}_3)_6[(\text{Cr}_{0.84}\text{S}_{0.16})\text{O}_4]_2 \cdot 12\text{H}_2\text{O}$
Monoclinic: $C2/c$
a 23.645, b 10.918, c 15.768 Å, β 114.42°
Pale yellow to bright lemon yellow; vitreous; transparent to translucent
Biaxial (+), α 1.647, β 1.674, γ 1.704,
2V(calc.) 88°
10.69 (100), 6.36 (50), 5.65 (50), 3.590 (70), 3.121 (80), 3.051 (80)
- IMA No. 96-050
The cadmium-dominant analogue of briartite
 $\text{Cu}_2\text{CdGeS}_4$
Tetragonal: $I42m$
a 5.45, c 10.6 Å

Colour unknown, only visible in polished section; metallic; opaque
 In reflected light: grey with pale violet tint, very weak anisotropism, very weak bireflectance and very weak pleochroism.
 R and ${}^{\text{im}}R$: (24.42, 10.79%) 460 nm, (23.29, 9.85%) 540 nm, (23.04, 9.59%) 580 nm, (23.46, 9.91%) 660 nm
 3.10 (100), 2.79 (10), 1.92 (80), 1.89 (70), 1.64 (60), 1.60 (20)

IMA No. 96-051

A polymorph of sibirskite
 $\text{Ca}_2\text{B}_2\text{O}_5 \cdot \text{H}_2\text{O}$
 Monoclinic: $P2_1/m$
 a 6.722, b 5.437, c 3.555 Å, β 93.00°
 White; weak pearly; translucent
 Biaxial (+), α 1.556, β 1.593, γ 1.663, $2V(\text{calc.})$ 75°
 6.73 (70), 3.354 (30), 2.975 (60), 2.855 (20), 2.237 (100), 1.776 (20)

IMA No. 96-052

The mercury-dominant analogue of černýite and stannite
 $\text{Cu}_2\text{HgSnS}_4$
 Tetragonal: $I\bar{4}$
 a 5.555, c 10.911 Å
 Dark grey; metallic; opaque
 In reflected light: greenish-grey to light grey with greenish-brownish tint, moderate anisotropism with faded colour effects form violet-blue to dark-greenish blue, insignificant bireflectance, weakly pleochroic from yellowish-olive-green to brownish-olive. $R_{\text{max.}}$: (26.0%) 470 nm, (26.3%) 546 nm, (25.6%) 589 nm, (24.8%) 650 nm
 3.17 (10), 1.958 (2.5), 1.941 (8), 1.671 (4), 1.646 (3.5), 1.264 (2.5)

IMA No. 96-053

The sulfate-dominant rhombohedral analogue of hydrocalumite
 $\text{Ca}_4\text{Al}_2(\text{OH})_{12}(\text{SO}_4) \cdot 6\text{H}_2\text{O}$
 Hexagonal (trigonal): $R\bar{3}$ or $R3$
 a 5.76, c 53.66 Å
 White; vitreous; transparent
 Uniaxial (–), ω 1.504, ε 1.485
 8.972 (100), 4.476 (70), 2.362 (40), 2.190 (40), 2.071 (35)

IMA No. 96-054

The Fe^{2+} -dominant analogue of hawthorneite
 $\text{Ba}[\text{Fe}_6^{2+}\text{Ti}_5\text{Mg}]\text{O}_{19}$
 Hexagonal: $P6_3/mmc$
 a 5.926, c 23.32 Å

Colour unknown, only visible in polished section; metallic; opaque
 In reflected light: light grey; very weak anisotropism, nearly isotropic; bireflectance very weak, but measurable; nonpleochroic.
 R_E , R_O ; ${}^{\text{im}}R_E$, ${}^{\text{im}}R_O$, $R_{\text{min.}}$: (16.9, 17.3; 5.13, 5.37%) 470 nm, (16.35, 16.8; 4.90, 5.19%) 546 nm, (16.3, 16.9; 4.92, 5.29%) 589 nm, (16.4, 17.1; 5.00, 5.42%) 650 nm
 2.963 (44), 2.795 (90), 2.641 (100), 2.437 (46), 1.676 (37), 1.634 (47), 1.481 (47)

IMA No. 96-055

The cerium-dominant analogue of 96-057, but structurally different
 $(\text{Ce,Nd,L a})\text{Al}(\text{SO}_4)_2(\text{C}_2\text{O}_4) \cdot 12\text{H}_2\text{O}$
 Monoclinic: $C2/c$
 a 8.718, b 18.313, c 13.128 Å, β 93.90°
 Very pale pink (incandescent light) and very pale blue (fluorescent light); vitreous; transparent
 Biaxial (+), α 1.455, β 1.485, γ 1.528, $2V(\text{meas.})$ 85°, $2V(\text{calc.})$ 82°
 7.9 (100), 5.36 (50), 5.01 (40), 3.93 (70), 3.74 (20), 3.29 (20), 3.07 (20)

IMA No. 96-056

$(\text{Ce,Nd,L a})_2(\text{SO}_4)_2(\text{C}_2\text{O}_4) \cdot 12\text{H}_2\text{O}$
 Triclinic: $P\bar{1}$
 a 6.007, b 8.368, c 9.189 Å, α 99.90°, β 105.55°, γ 107.71°
 Pale pink (incandescent light), pale blue (fluorescent light), some cream-coloured; vitreous; transparent
 Biaxial (–), α 1.544, β 1.578, γ 1.602, $2V(\text{meas.})$ 65°, $2V(\text{calc.})$ 78°
 8.52 (70), 6.72 (60), 5.48 (100), 4.26 (50), 3.84 (60), 3.35 (40), 2.744 (40)

IMA No. 96-057

The yttrium-dominant analogue of 96-055, but structurally different
 $(\text{Y,Nd,Ce})\text{Al}(\text{SO}_4)_2(\text{C}_2\text{O}_4) \cdot 12\text{H}_2\text{O}$
 Monoclinic: $P2/n$
 a 10.289, b 19.234, c 11.015 Å, β 108.50°
 Colourless; vitreous; transparent
 Biaxial (+), α 1.48, β 1.49, γ 1.55, $2V(\text{meas.})$ 7°, $2V(\text{calc.})$ 46°
 9.3 (100), 6.28 (90), 5.20 (40), 4.89 (60), 4.63 (30), 4.09 (50), 3.700 (30)

IMA No. 96-058

The rubidium-dominant analogue of microcline
 $(\text{Rb,K})\text{AlSi}_3\text{O}_8$
 Triclinic: $P\bar{1}$

a 8.81, b 13.01, c 7.18 Å, α 90.3°, β 115.7°, γ 88.2°

Colourless; vitreous; transparent

Biaxial, indices of refraction slightly higher than host microcline

5.82, 5.77, 4.62, 3.88, 3.61, 3.60, 3.59, 2.94, 2.65, 2.63, 2.61, 2.56 (electron diffraction, no intensities)

IMA No. 96-059

$\text{Fe}^{3+}\text{Mo}_2\text{O}_6(\text{OH})_3 \cdot \text{H}_2\text{O}$

Triclinic: $\text{P}\bar{1}$

a 5.878, b 7.536, c 9.436 Å, α 71.66°, β 83.43°, γ 72.85°

Green with a yellowish tinge; vitreous to earthy; transparent to opaque

Biaxial (-), α 1.91, β 2.03, γ 2.11, $2V(\text{meas.}) \sim 90^\circ$, $2V(\text{calc.}) 74^\circ$

5.620 (70), 4.711 (50), 4.095 (70), 3.319 (100), 3.232 (90), 2.614 (50), 1.956 (50)

IMA No. 96-060

The scandium-dominant analogue of overite and segelerite

$\text{CaMgSc}(\text{PO}_4)_2(\text{OH}) \cdot 4\text{H}_2\text{O}$

Orthorhombic: Pbca

a 15.03, b 18.95, c 7.59 Å

Colourless, light yellow to yellowish brown; vitreous; transparent

Biaxial (-), α 1.574, β 1.579, γ 1.582, $2V(\text{meas.}) \sim 50^\circ$, $2V(\text{calc.}) 75^\circ$

9.49 (100), 4.75 (17), 3.440 (31), 2.942 (27), 2.912 (44), 2.890 (35), 2.018 (15)

IMA No. 96-062

$(\text{Ti,Cr,Fe})[\text{O}_{2-x}(\text{OH})_x]$

Monoclinic: $\text{P}2_1/c$

a 7.688, b 4.5495, c 20.147 Å, β 92.27°

Black; metallic; translucent to opaque

Biaxial, mean n 2.47 (calc.). In reflected light: grey, with R lower than that of rutile, crichtonite, and srilankite and higher than that of pyrope

3.766 (66), 2.835 (100), 2.660 (73), 1.6842 (94), 1.6760 (73), 1.6574 (71)

IMA No. 96-063

The sodium-dominant analogue of lemoynite with additional H_2O

$\text{Na}_4\text{Zr}_2\text{Si}_{10}\text{O}_{26} \cdot 9\text{H}_2\text{O}$

Monoclinic: $\text{C}2/m$

a 10.5150, b 16.2534, c 9.1029 Å, β 105.46°

Colourless to white; vitreous; transparent to translucent

Biaxial (-), α 1.533, β 1.559, γ 1.567, $2V(\text{meas.}) 63^\circ$, $2V(\text{calc.}) 57^\circ$

8.832 (30), 8.135 (100), 5.975 (40), 3.974 (35), 3.693 (30), 3.564 (40), 3.490 (35), 2.804 (30)