Zeitschrift:	Schweizerische mineralogische und petrographische Mitteilungen = Bulletin suisse de minéralogie et pétrographie
Band:	80 (2000)
Heft:	3
Artikel:	U-Pb zircon dating of two contrasting Late Variscan plutonic suites from the Pelvoux massif (French Western Alps)
Autor:	Guerrot, Catherine / Debon, François
DOI:	https://doi.org/10.5169/seals-60964

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. <u>Siehe Rechtliche Hinweise.</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. <u>Voir Informations légales.</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. <u>See Legal notice.</u>

Download PDF: 29.04.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

U-Pb zircon dating of two contrasting Late Variscan plutonic suites from the Pelvoux massif (French Western Alps)

by Catherine Guerrot¹ and François Debon²

Abstract

The Pelvoux massif comprises extensive Late Variscan plutons emplaced into a polymetamorphic and migmatitic basement. On the basis of their average Mg / (Fe + Mg) ratios, the plutons can be divided into a magnesian and a magnesian-ferriferous suite. Age determinations by the U–Pb conventional technique on zircon on a representative pluton of each suite gave 343 ± 11 Ma (Viséan) for the magnesian Rochail pluton and 302 ± 2 Ma (Stephanian) for the magnesian-ferriferous Turbat-Lauranoure pluton. By comparison with other External Crystalline Massifs of the Alps, these ages are interpreted to indicate that the generation of the Late Variscan plutons in the Pelvoux massif was a discontinuous process, involving two major magmatic pulses separated by a gap of some 40 Ma.

Keywords: plutonic rocks, geochemistry, geochronology, Variscan orogeny, Pelvoux massif, Western Alps.

1. Introduction

Late Variscan plutonic rocks are widely represented within the External Crystalline Massifs of the Alps or ECM (Fig. 1; DEBON and LEMMET, 1999, and references therein). But for this plutonism to be understood requires, among other things, meaningful age determinations based upon methods little sensitive to tectono-metamorphic processes (e.g. SCHALTEGGER, 1994). The U-Pb zircon datings we present here fulfil this requirement. They concern two plutons of the Pelvoux massif, viz. a domain where reliable intrusion ages were almost completely lacking. On the basis of geochemical data, two Late Variscan plutonic suites can be distinguished in this massif, namely a highly magnesian (high-mg-number) suite and a more iron-rich magnesian-ferriferous (low-mg-number) suite (DEBON and LEMMET, 1999; Figs 2 and 3). The aim of our paper is to present the data which substantiate the age of emplacement of the two suites and to situate the latter within the frame of the Late Variscan plutonic history of the ECM.

2. Dated plutonic bodies

More than twenty Late Variscan plutons occur in the Pelvoux massif, covering about one third of its area (Fig. 2). Based on data from the literature (e.g. LE FORT, 1973; BARFÉTY and PÊCHER, 1984), their main characteristics are summarized in DEBON and LEMMET (1999) who divided them into two suites of different *mg*-number. A typical representative of each suite was selected for U– Pb zircon dating, namely the magnesian Rochail pluton and the magnesian-ferriferous Turbat-Lauranoure pluton (Figs 2 and 3).

2.1. ROCHAIL PLUTON

The Rochail pluton (ca. 34 km^2) intrudes gneissic formations and comprises two granitic units (BOISSET, 1986; BARFÉTY, 1989). The main unit, making up the southern part of the pluton, is composed of a massive to weakly foliated, mediumgrained monzogranite with biotite \pm rare muscovite, apatite and zircon. To the north, this merges

¹ BRGM, B.P. 6009, F-45060 Orléans Cedex 02, France. <c.guerrot@brgm.fr>

² LGCA, UMR 5025 CNRS, Grenoble, France. Present address: 3, Chemin du Magnit, F-38490 Le Passage, France.



Fig. 1 External Crystalline Massifs of the Alps (modified from DEBON and LEMMET, 1999). AA, Aar AG, Argentera AR, Aiguilles Rouges BE, Belledonne GO, Gotthard GR, Grandes Rousses MB, Mont Blanc PE, Pelvoux (study area).



Fig. 2 Geological map of the Pelvoux massif (after DEBON and LEMMET, 1999). Typology of the Variscan plutons is based on their plots on the *mg*-number-B diagram (see figure 3).



Fig. 3 Plot of the Pelvoux plutons (average compositions) on the *mg*-number-B diagram (after DEBON and LEMMET, 1999). Reference system of mean compositions: gr, granite; ad, adamellite; gd, granodiorite; to, to-nalite; qd, quartz diorite. For numbering of plutons see figure 2.

into a coarser and darker foliated biotite monzogranite with a porphyritic tendency. Both monzogranites enclose conspicuous mafic igneous enclaves, especially of the vaugneritic-durbachitic type (the so-called "syénite du Lauvitel").

The emplacement age of the Rochail pluton has remained poorly constrained despite a number of radiometric studies performed both on granites and vaugneritic enclaves. Besides an age of 331 ± 32 Ma (2σ) ascribed to the northern unit from a questionable Rb-Sr whole-rock isochron (initial 87 Sr/ 86 Sr ratio of 0.7047 ± 0.0013) (DE-MEULEMEESTER, 1982), a variety of ages had been obtained through the total-lead zircon method (271, 298, 308 Ma; CHESSEX et al., 1964; KRUM-MENACHER et al., 1965) and Rb-Sr or K-Ar techniques applied to both biotite (fourteen ages ranging from 267 to 339 Ma) and amphibole (five ages from 321 to 378 Ma) (BONHOMME et al., 1963; DEMEULEMEESTER, 1982; VITTOZ et al., 1987; G. BANZET, pers. commun.).

The dated sample (RO 1; Fig. 2; Tab. 1) was collected in situ on the northwestern side of Lake Lauvitel ($x = 6^{\circ}03'41"$, $y = 44^{\circ}58'28"$, z = 1595 m) from a homogeneous, light-coloured, massive, medium-grained, quartz-bearing syenite with biotite and rare muscovite, about one metre away from a vaugneritic enclave more than $5 \times 2 m$ in size. This syenite, quite different from the so-called "syénite du Lauvitel", merges laterally into the dominant monzogranitic type, of which it represents a quartz-poor and feldspar-rich variety.



Fig. 4 Distribution of zircons from the dated samples (RO 1, TL 1) in the typological diagram of PUPIN (1980).

Pluton	Rochail	Turbat-Lauranoure
Dated sample	RO 1	TL 1
SiO	63.49	71.85
TiO	0.32	0.26
AlaOa	19.40	14 35
Fe ₂ O ₂ *	1.60	2 30
MnO	tr	0.03
MgO	1.03	0.56
CaO	0.92	1.61
Na ₂ O	5.48	3.17
K ₂ Õ	6.18	4.74
P_2O_5	0.13	0.13
LOI	1.09	0.78
Total	99.65	99.78
B = Fe + Mg + Ti	50	46
Mg/(Fe + Mg)	0.56	0.33
Ba	1498	563
Cr	25.7	6.7
Hf	6.8	4.8
Nb	6.9	10.8
Ni	12.9	3.1
Rb	244	213
Sr	814	137
Ta	1.8	1.6
Th	61.6	16.0
	12.3	3.0
	26.3	20.2
VV V	0.6	0.0
	3./ 210	150
	219	138
La	10.14	32.35
Ce	19.64	67.07
Pr	2.21	7.58
Nd	8.58	28.87
Sm	1.61	6.14
Eu	0.81	0.80
Ga	1.22	4.78
	0.16	0.75
	0.90	4.15
Er	0.21	2 10
Tm	0.37	0.35
Vh	0.10	0.55
	0.05	0.37
Lu	0.12	0.57

Major elements in oxide-%; trace elements in ppm; $Fe_2O_3^*$, total iron as ferric oxide; LOI, loss on ignition; tr, traces. Analyses performed at the CRPG-CNRS (Nancy), by ICP-Emission (major elements) and ICP-MS (trace elements).

2.2. TURBAT-LAURANOURE PLUTON

The Turbat-Lauranoure pluton (ca. 33 km²) intrudes gneissic and migmatitic formations (LE FORT, 1973; BARFÉTY and PÊCHER, 1984). It is a homogeneous body of variably porphyritic and foliated, coarse-grained biotite \pm muscovite syenogranite enclosing rare microgranular enclaves. No previous radiometric age is available for this pluton, apart from a questionable whole-rock Rb–Sr age of ca. 300 Ma (310 Ma with new constants) given by three samples collected from three magnesian-ferriferous plutons (*i.e.* Etages, Orgières, Turbat-Lauranoure; J. SONET, in LE FORT, 1973).

The dated sample (TL 1) was collected in situ on the northern side of the Bonne river ($x = 6^{\circ}08'46''$, $y = 44^{\circ}52'57''$, z = 1590 m) from a massive syenogranite; it is representative of the main rock type (Fig. 2; Tab. 1).

3. U–Pb zircon geochronology

3.1. ANALYTICAL PROCEDURE

After crushing the sample and concentrating the heavy minerals with heavy liquids, the zircons were separated into fractions of different magnetic susceptibility, non-magnetic zircons being generally more concordant (KROGH, 1982a). Zircons from the least magnetic fraction were examined under a binocular microscope, and some 30 to 50 grains selected according to optical and morphological criteria. These were then air-abraded in order to remove their outer surfaces so as to minimize effects due to lead loss (KROGH 1982b). Techniques of zircon dissolution and chemical separation of Pb and U were adapted from KROGH (1973) and PARRISH (1987). All samples were spiked using a mixed ²⁰⁵Pb-²³³U-²³⁵U isotope tracer added before dissolution. Pb and U were loaded on the same single Re filament with silica-gel and H₃PO₄ and analyses were performed on a Finnigan MAT 261 mass spectrometer in single collector operation, using a secondary electron multiplier. Mass discrimination, controlled by regular measurements of the NBS-981 common Pb standards, was 0.28% / a.m.u. Total procedure blanks were less than 15 pg for Pb and 1 pg for U. Pb analyses were corrected for a blank isotopic composition of ${}^{206}Pb/{}^{204}Pb = 18.0$, ${}^{207}Pb/{}^{204}Pb = 18.0$, ${}^{207}Pb/{}^{204}Pb = 18.0$ 206 Pb = 0.85, 208 Pb/ 206 Pb = 2.06. Individual errors are given at 2σ . Regressions were calculated according to DAVIS (1982), and also according to LUDWIG (1999) for lower intercept and concordia age. Quoted errors are given at the 95% confidence level.

3.2. ROCHAIL PLUTON

The analysed sample from the Rochail pluton (RO 1) contains abundant pale-pink euhedral zir-

Tab. 1 Chemical composition of the dated samples (RO 1, TL 1).

	Sample		Conc	centratio	SU		Is	otopic Ratios			A	pparent Age	
3				1	Pb	;	;			ŝ			Ē
#	Grain	weight	D	Pb rad	com	Pb	Pb	Pb/U	Pb/U	Чb	Pb/U	Pb/U	PD 01
		(au)	(mdd)	(mdd)	(Dg)	206/204	208/206	206/238	207/235	207/206	206/238	207/235	207/206
	[1]				2	[3]	[4]	[4]	[4]	[4]	(Ma)	(Ma)	(Ma)
RO	CHAIL PLU	TON R	01										
A	3.c.sp.eu.m1	12	431.1	21.8	2	2317	0.21	0.0518 ± 4	0.4788 ± 34	0.06701 ± 18	325.7	397.2	838.0
В	1.p.lp.eu.m1	6	241.6	12.5	18	370	0.52	0.0484 ± 2	0.3559 ± 31	0.05332 ± 40	304.8	309.2	342.6
C	5.p.lp.eu.ml	27	1251.3	75.4	293	440	0.38	0.0597 ± 3	0.5975 ± 33	0.07263 ± 22	373.6	475.6	1003.6
Ω	1.p.sp.sb.ml	23	1142.7	58.6	122	682	0.38	0.0500 ± 2	0.3679 ± 20	0.05332 ± 26	314.7	318.1	342.6
Ц	1.c.t.m1	S	1075.1	53.6	62	295	0.16	0.0525 ± 2	0.3859 ± 24	0.05332 ± 24	329.8	331.4	342.6
IJ	1,p,lp,eu,m1	2	825.9	168.0	16	1227	0.83	0.1979 ± 10	3.3283 ± 180	0.12199 ± 32	1163.9	1487.7	1985.5
UT	RBAT-LAUR	ANOL	IRE PLU	TONT	L1								
A	3.p.sp.eu.m0	13	753.4	69.69	24	2373	0.06	0.0935 ± 8	1.1468 ± 120	0.08901 ± 60	575.9	775.7	1404.3
C	5.p.lp.eu.m0	52	442.4	25.7	82	1048	0.08	0.0594 ± 3	0.4573 ± 26	0.05580 ± 26	372.2	382.4	444.4
Ω	4,c.sp.eu.m0	35	456.2	21.5	66	491	0.09	0.0480 ± 3	0.3465 ± 32	0.05240 ± 40	302.0	302.1	303.0
Ш	1.p.lp.sb.m1	4	846.7	43.4	280	57	0.10	0.0514 ± 12	0.4184 ± 243	0.05903 ± 346	323.2	354.9	568.2
ſĿ,	1,c,lp,eu,m1	4	808.5	40.6	27	388	0.09	0.0507 ± 3	0.3958 ± 58	0.05658 ± 78	319.1	338.6	475.0
G	1,v,sp,eu,m1	4	1443.0	74.1	29	681	0.04	0.0548 ± 5	0.4118 ± 60	0.05445 ± 70	344.2	350.2	389.9
[1] =	number of gr	ains, c c	solourles	s, p pink	, v viol	et, lp long	g prism, sl	p short prism, eu	ı euhedral, sb subhe	dral, t tip, m0 magneti	ic at 0° tilt, m1	_	

U–Pb analytical results for zircons from the dated samples (RO 1, TL 1)

Tab. 2

Pb (spike, contamination and mineral) = Total common magnetic at 1° till

= corrected for mass fractionation = corrected for mass fractionation,

Pb according to STACEY and KRAMERS (1975) blank (Pb=15 pg, U=1 pg), spike, and initial common Uncertainties are given at $\pm 2\sigma$ and refer to the last decimals 004

cons, commonly asymmetric or displaying internal zoning. Most of the crystals show small black inclusions. One hundred crystals were indexed; this population, scattered near the centre of the classification diagram of PUPIN (1980), has I.A, I.T mean values of 388 and 386 respectively (Fig. 4).

Six zircon fractions (each of 1 to 5 individual grains) were analysed from this sample. Three fractions (B, D and F, Tab. 2) are collinear and show a degree of discordance from 3 to 10%, reflecting lead loss. Despite rather low ²⁰⁶Pb/²⁰⁴Pb ratios probably related to the presence of small inclusions, they define a discordia intersecting the concordia at 343 +11/-10 Ma (92% probability of fit) (Fig. 5), the lower unforced intercept being at the origin of the diagram. The other three fractions (especially fraction G) reveal the presence of an old inherited component, probably of Early Proterozoic age. No alignment is seen from these three fractions, probably due to some combination of inheritance and lead loss.

The age of 343 +11/-10 Ma, i.e. Early Viséan (ODIN, 1994), is interpreted as the intrusion age of the Rochail pluton. Because of their field relationships, it is most likely that this age can be ascribed also to vaugneritic-durbachitic the enclaves.

3.3. TURBAT-LAURANOURE PLUTON

The sample from the Turbat-Lauranoure pluton (TL 1) contains rather abundant transparent pink zircon crystals. One hundred crystals were indexed; this population, widely scattered in the centre of PUPIN's (1980) classification diagram, has I.A, I.T mean values of 500 and 540 respectively (Fig. 4), comparable with those of 575 and 429 previously obtained by COSTARELLA (1987).

Of the six fractions analysed, only fraction D is concordant at 302 ± 2 Ma. The other fractions show









evidence of inherited lead (Tab. 2). Fraction E, with a high common lead content, was not considered for further calculation. Two alignments are observed in the Concordia diagram (Fig. 6). The first, with fractions D, C and G, defines a regression with an upper intercept at 692 ± 66 Ma and a lower intercept at 302 \pm 11 Ma (68% probability of fit). The second alignment (D, F, A) gives an upper intercept at 1913 ± 22 Ma and a lower intercept at 302 ± 2 Ma (95% probability of fit). Fraction E falls on this regression line. This sample thus shows evidence of Cadomian and Lower Proterozoic inheritance. In relation with the concordant position of fraction D, and with the two linear alignments obtained, a weighted average mean age of 302 ± 2 Ma, i.e. Stephanian (ODIN, 1994), can be assigned to the crystallization of the Turbat-Lauranoure pluton.

4. Conclusion

Apart from an unpublished age of 312 ± 7 Ma ascribed to the magnesian Combeynot granite (Fig. 2) by CANNIC et al. (1998; see also DEBON and LEMMET, 1999), this study provides the first reliable and reasonably precise U–Pb zircon ages on plutonic rocks from the Pelvoux massif, which is of major interest for the reconstitution of the magmatic evolution of the Variscan belt in the External Crystalline Massifs of the Alps.

The Late Variscan plutonic bodies, in the ECM as a whole, can be divided into an early (Viséan; ca. 330–340 Ma) high-mg-number suite, and a later (mainly Stephanian; ca. 295–305 Ma) low-mgnumber suite (DEBON and LEMMET, 1999). This two-fold partition is, in all respects (typology, age), quite comparable with that defined in the Pelvoux massif. It is likely that the Late Variscan plutonic activity in the latter was comparable to that evidenced elsewhere in the ECM, viz. a discontinuous process comprising two major events separated by a gap of up to some 40 Ma. The first event would include most of the magnesian plutons, followed by a second event with more ironrich, magnesian-ferriferous plutons.

Acknowledgements

Early drafts of the manuscript were greatly improved by critical comments from F. Bussy (University of Lausanne) and U. Schaltegger (ETH, Zürich). The authors gratefully acknowledge G. Banzet (CNRS), J.C. Barféty (BRGM), P. Jézéquel (BRGM), P. Le Fort (CNRS), A. Pêcher (University of Grenoble), M. Robert and D. Thiéblemont (BRGM) for helpful advice or technical assistance, and M. Engi for careful editorial handling.

References

- BARFÉTY, J.C. (Coord.) (1989): Carte géologique de la France à 1/50 000, feuille La Mure (821), et notice explicative. BRGM, Orléans, 98 pp.
- BARFÉTY, J.C. and PÊCHER, A. (Coords) (1984): Carte géologique de la France à 1/50 000, feuille Saint-Christophe-en-Oisans (822), et notice explicative. BRGM, Orléans, 64 pp.
- BOISSET, T. DE (1986): Les enclaves basiques du granite du Rochail (syénite du Lauvitel), Haut Dauphiné, Alpes françaises. Pétrographie et minéralogie. Thèse de 3^{ème} Cycle, Univ. Grenoble, 188 pp.
- BONHOMME, M., BUFFIÈRE, J.M., MICHEL, R., PHILIBERT, J., ROQUES, M. and VIALETTE, Y. (1963): Age absolu du massif du Rochail (secteur Nord-Ouest du massif du Pelvoux, Isère). C.R. Acad. Sci. Paris 256, 2649– 2651.
- CANNIC, S., LAPIERRE, H., SCHÄRER, U., MONIÉ, P., BRI-OUEU, L. and BASILE, C. (1998): Origin of Hercynian magmatism in the French Western Alps: geochemical and geochronological constraints. Abstract (Goldschmidt Conf., Toulouse, 1998). Mineral. Mag. 62A, 274–275.
- CHESSEX, R., DELALOYE, M., KRUMMENACHER, D. and VUAGNAT, M. (1964): Nouvelles déterminations d'âges "plomb total"sur des zircons alpins. Schweiz. Mineral. Petrogr. Mitt. 44, 43–60.
- COSTARELLA, R. (1987): Le complexe annulaire alcalin de Combeynot (Massifs Cristallins Externes, Alpes françaises), témoin d'un magmatisme en régime distensif. Pétrogéochimie et signification géodynamique. Thèse 3ème Cycle, Univ. Grenoble, 268 pp.
- DAVIS, D.W. (1982): Optimum linear regression and error estimation applied to U–Pb data. Can. J. Earth Sci. 19, 2141–2149.
- DEBON, F. and LEMMET, M. (1999): Evolution of Mg/Fe ratios in Late Variscan plutonic rocks from the External Crystalline Massifs of the Alps (France, Italy, Switzerland). J. Petrol. 40, 1151–1185.
- DEMEULEMEESTER, P. (1982): Contribution à l'étude radiométrique à l'argon et au strontium des massifs cristallins externes (Alpes françaises). Distribution cartographique des âges sur biotites et amphiboles. Thèse 3^{ème} Cycle, Univ. Grenoble, 227 pp.
- KROGH, T.E. (1973): A low-contamination method for hydrothermal decomposition of zircon and extraction of U and Pb for isotopic age determination. Geochim. Cosmochim. Acta 37, 485–494.
- KROGH, T.E. (1982a): Improved accuracy of U–Pb zircon dating by selection of more concordant fractions using a high gradient magnetic separation technique. Geochim. Cosmochim. Acta 46, 631–635.
- KROGH, T.E. (1982b): Improved accuracy of U–Pb zircon ages by the creation of more concordant systems using an air abrasion technique. Geochim. Cosmochim. Acta 46, 637–649.
- KRUMMENACHER, D., BORDET, P. and LE FORT, P. (1965): Les massifs externes alpins et leurs séries métamorphiques. Problèmes de raccords stratigraphiques et géochronométriques. Schweiz. Mineral. Petrogr. Mitt. 45, 855–874.
- LE FORT, P. (1973): Géologie du Haut-Dauphiné cristallin (Alpes françaises). Etude pétrologique et structurale de la partie occidentale. Mém. Sci. Terre 25, 373 pp.
- LUDWIG, K.R. (1999): Isoplot/Ex version 2.06: A Geochronological Toolkit for Microsoft Excel. Berkeley Geochronology Center, Special Publication N° 1a, 49 pp.

- ODIN, G.S. (1994): Geological time scale. C.R. Acad. Sci. Paris 318 II, 59–71.
- PARRISH, R.R. (1987): An improved micro-capsule for zircon dissolution in U–Pb geochronology. Chem. Geol. (Isotop. Geosci. Sect.) 66, 99–102.
- Geol. (Isotop. Geosci. Sect.) 66, 99–102.
 PUPIN, J.P. (1980): Zircon and granite petrology. Contrib. Mineral. Petrol. 73, 207–220.
 SCHALTEGGER, U. (1994): Unravelling the pre-Mesozoic
- SCHALTEGGER, U. (1994): Unravelling the pre-Mesozoic history of Aar and Gotthard massifs (Central Alps) by isotopic dating – a review. Schweiz. Mineral. Petrogr. Mitt. 74, 41–51.
- VITTOZ, P., BONHOMME, M.G. and VIVER, G. (1987): Datation K/Ar des minéraux des rohes ignées de la région du Lauvitel (Haut-Dauphiré, France). Géol. Alpine 63, 165–172.

Manuscript received November D, 1999; revision accepted September 21, 2000.